



AMERICAN  
COLLEGE *of*  
CARDIOLOGY®

## ACC Clinical Bulletin

# Cardiac Implications of Novel Wuhan Coronavirus (COVID-19)

### Background on Coronavirus epidemic <sup>i,ii</sup>

- COVID-19 was first reported in late December 2019, originating in Wuhan, China
- COVID-19 is a betacoronavirus, like SARS and MERS, presenting as viral pneumonia with a wide range of acuity
- As of February 12, there are 45,204 confirmed cases and 1,117 confirmed deaths across 28 countries; COVID-19 appears to have greater infectivity and a lower case fatality rate when compared to SARS and MERS
- 99% of all cases are in mainland China, where despite aggressive containment efforts, case counts continue to rise rapidly

### Early cardiac implications from case reports on Wuhan Coronavirus<sup>iii</sup>

- Early case reports suggest patients with underlying conditions are at higher risk for complications or mortality from COVID-19; up to 50% of hospitalized patients have a chronic medical illness
- 40% of hospitalized patients with confirmed COVID-19 patients have cardiovascular or cerebrovascular disease
- In a recent case report on 138 hospitalized COVID-19 patients, 19.6% of patients developed acute respiratory distress syndrome<sup>iv</sup>
  - 16.7% of patients developed arrhythmia; 7.2% developed acute cardiac injury
  - 8.7% of patients developed shock; 3.6% developed acute kidney injury
  - Rates of complication were universally higher for ICU patients



- The first reported death was a 61-year-old male, with a long history of smoking, who succumbed to acute respiratory distress, heart failure, and cardiac arrest
- Early, unpublished first-hand reports suggest at least some patients develop myocarditis

### Potential cardiac implications from analog viral respiratory pandemics

- **Influenza analog:** In all influenza pandemics other than the 1918 flu, cardiovascular events surpassed all other causes of mortality, including superimposed pneumonia<sup>v</sup>
- **General viral analog:** Viral illness is a well-known destabilizing factor in chronic cardiovascular disease, a general consequence of the imbalance between infection-induced increased metabolic demand and reduced cardiac reserve. The viral infection along with superimposed pneumonia will directly and indirectly affect the cardiovascular system<sup>vi</sup>
  - Both coronary artery disease and heart failure patients are at increased risk of acute events or exacerbation; viral illness can potentially destabilize coronary plaques through several mechanisms including systemic inflammatory responses which have been recently documented with COVID-19
  - Multiple co-morbidities (DM, obesity, HTN, COPD, CKD) further increase risk
- **SARS/MERS analog:** Although published literature on CV implications of SARS/MERS is limited, in the absence of more detailed reporting on COVID-19, it may prove instructive
  - 60% of MERS cases had one or more pre-existing comorbidity, resulting in a poorer prognosis; expert guidance suggests patients with diabetes, CVD, or renal disease should be prioritized for treatment<sup>vii</sup>
  - Both SARS and MERS have been linked to acute myocarditis, acute myocardial infarction, and rapid-onset heart failure
    - In one early published report, 2 out of the 5 deaths were attributed to MI<sup>viii,ix</sup>
    - These data should be interpreted cautiously—indicative of the increased CV risk in coronavirus patients, but not generalizable to broader outcomes
  - Reversible, sub-clinical diastolic LV impairment in acute SARS even among those without underlying cardiac disease appears common, likely the result of systemic inflammatory immune response and is not unique to SARS; however, lower EF upon admission was predictive of later mechanical ventilation<sup>x</sup>



**ACC Clinical Bulletin**  
Cardiac Implications  
of Novel Wuhan Coronavirus  
(COVID-19)

- In one study of cardiovascular complications of SARS in 121 patients:<sup>xi</sup>
  - 71.9% of patients developed persistent tachycardia, including 40% with continued tachycardia during outpatient follow-up
  - 50.4% of patients developed sustained asymptomatic hypotension during hospitalization; one patient required inotropic support
  - 14.9% of patients developed transient bradycardia
  - 10.7% of patients developed transient cardiomegaly, without signs or symptoms of heart failure
  - One patient experienced transient paroxysmal AF, with spontaneous resolution
  - Cardiovascular complications appeared statistically uncorrelated with oxygen desaturation or ICU admission

**Clinical guidance given current COVID-19 uncertainty**

- COVID-19 is spread through droplets and can live for substantial periods outside the body; containment and prevention using standard public health and personal strategies for preventing the spread of communicable disease remains the priority
- In geographies with active COVID-19 transmission (mainly China), it is reasonable to advise patients with underlying cardiovascular disease of the potential increased risk and to encourage additional, reasonable precautions
- Older adults are less likely to present with fever, thus close assessment for other symptoms such as cough or shortness of breath is warranted
- Some experts have suggested that the rigorous use of guideline-directed, plaque stabilizing agents could offer additional protection to CVD patients during a widespread outbreak (statins, beta blockers, ACE inhibitors, ASA)<sup>v</sup>; however, such therapies should be tailored to individual patients
- It is important for patients with CVD to remain current with vaccinations, including the pneumococcal vaccine given the increased risk of secondary bacterial infection; it would also be prudent to receive influenza vaccination to prevent another source of fever which could be initially confused with coronavirus infection
- It may be reasonable to triage COVID-19 patients according to the presence of underlying cardiovascular, respiratory, renal, and other chronic diseases for prioritized treatment
- Providers are cautioned that classic symptoms and presentation of AMI may be overshadowed in the context of coronavirus, resulting in underdiagnosis



**ACC Clinical Bulletin**  
Cardiac Implications  
of Novel Wuhan Coronavirus  
(COVID-19)

- For CVD patients in geographies without widespread COVID-19 emphasis should remain on the threat from influenza, the importance of vaccination and frequent handwashing, and continued adherence to all guideline-directed therapy for underlying chronic conditions
- COVID-19 is a fast-moving epidemic with an uncertain clinical profile; providers should be prepared for guidance to shift as more information becomes available

- 2019 Novel Coronavirus Situation Summary, Centers for Disease Control (February 10, 2020), retrieved from <https://www.cdc.gov/coronavirus/2019-nCoV/summary.html>
- Coronavirus COVID-19 Global Cases by Johns Hopkins CSSE (February 12, 2020), retrieved from [https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html?fbclid=IwAR29qGU1Zs2huweaWHXJA7sl\\_YnkdDNreGxKeH7qMHVVqXvuymQVB-DrSBg#/bda7594740fd40299423467b48e9ecf6](https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html?fbclid=IwAR29qGU1Zs2huweaWHXJA7sl_YnkdDNreGxKeH7qMHVVqXvuymQVB-DrSBg#/bda7594740fd40299423467b48e9ecf6)
- Chen H, Zhou M, Dong X, et al. Epidemiological and Clinical Characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020; published online January 29. <https://www.thelancet.com/action/showPdf?pii=S0140-6736%2820%2930211-7>
- Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients with 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. Published online February 07, 2020. doi:10.1001/jama.2020.1585
- Madjid M, Casscells SW. Of birds and men: cardiologists' role in influenza pandemics. *Lancet* 2004. 364: 1309.
- Expert opinion
- Al-Abdely, HM, Midgley, CM, Alkhamis AM, et al. Middle East Respiratory Syndrome Coronavirus Infection Dynamics and Antibody Responses among Clinically Diverse Patients, Saudi Arabia. *Emerging Infectious Diseases* 2019. 25(4): 753-766.
- Alhagbani T. Acute myocarditis associated with novel Middle East respiratory syndrome coronavirus. *Annals of Saudi Medicine* 2016. 36(1): 78-80.
- Peiris JM, Chu CM, Cheng VC. Clinical Progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet* 2003. Retrieved from <http://image.thelancet.com/extras/03art4432web.pdf>.
- Li SS, Cheng C, Fu C, et al. Left Ventricular Performance in Patients with Severe Acute Respiratory Syndrome: A 30-Day Echocardiographic Follow-Up Study. *Circulation* 2003. 108: r93-98).
- Yu CM, Wong R, Wu EB. Cardiovascular complications of severe acute respiratory syndrome. *Postgrad Med J* 2006. 82:140-144.

## Expert Advisors

### Mohammad Madjid, MD, MS, FACC, FSCAI

Assistant Professor of Medicine  
McGovern Medical School  
University of Texas at Houston Health Science Center  
Memorial Hermann Heart and Vascular Institute

### Scott D Solomon, MD, FACC

The Edward D. Frohlich Distinguished Chair  
Professor of Medicine  
Harvard Medical School  
Brigham and Women's Hospital

### Orly Vardeny, PharmD, MS, FAHA, FHFSA

Minneapolis VA Center for Care Delivery and Outcomes Research  
Associate Professor of Medicine  
University of Minnesota

## Correspondence

### Brendan Mullen, EVP, Science & Quality, American College of Cardiology

[bmullen@acc.org](mailto:bmullen@acc.org)

## Reviewed and Approved

February 12, 2020 ACC Science and Quality Oversight Committee