

**Title: Management of Myocardial Revascularization Failure: An Expert Consensus Document of the EAPCI.**

**Authors:** Giulio G. Stefanini, M.D, PhD; Fernando Alfonso, M.D, PhD; Emanuele Barbato, M.D, PhD; Robert A. Byrne, MB, BCh, PhD; Davide Capodanno, M.D, PhD; Roisin Colleran, MB, BCh; Javier Escaned, M.D, PhD; Daniele Giacoppo, M.D; Vijay Kunadian, MBBS, PhD; Alexandra Lansky, M.D; Julinda Mehilli, M.D; Franz-Josef Neumann, M.D; Damiano Regazzoli, M.D; Jorge Sanz-Sánchez, M.D, PhD; William Wijns, M.D, PhD; Andreas Baumbach, M.D, PhD.

**DOI:** 10.4244/EIJ-D-20-00487

**Citation:** Stefanini GG, Alfonso F, Barbato E, Byrne RA, Capodanno D, Colleran R, Escaned J, Giacoppo D, Kunadian V, Lansky A, Mehilli J, Neumann FJ, Regazzoli D, Sanz-Sánchez J, Wijns W, Baumbach A. Management of Myocardial Revascularization Failure: An Expert Consensus Document of the EAPCI. *EuroIntervention* 2020; Jaa- 2020, doi: 10.4244/ EIJ-D-20-00487

**Guest Editor:** Alec Vahanian, M.D, PhD

**Manuscript submission date:** 23 April 2020

**Revisions received:** 01 June 2020, 15 June 2020

**Accepted date:** 25 June 2020

**Online publication date:** 30 June 2020

**Disclaimer:** This is a PDF file of a "Just accepted article". This PDF has been published online early without copy editing/typesetting as a service to the Journal's readership (having early access to this data). Copy editing/typesetting will commence shortly. Unforeseen errors may arise during the proofing process and as such Europa Digital & Publishing exercise their legal rights concerning these potential circumstances.

# MANAGEMENT OF MYOCARDIAL REVASCULARIZATION FAILURE: AN EXPERT CONSENSUS DOCUMENT OF THE EAPCI

## *Short title: EAPCI Consensus Document on the Management of Myocardial Revascularization Failure*

Giulio G. Stefanini, MD, PhD<sup>1,2\*</sup>; Fernando Alfonso, MD, PhD<sup>3</sup>; Emanuele Barbato, MD, PhD<sup>4</sup>; Robert A. Byrne, MB, BCh, PhD<sup>5,6</sup>; Davide Capodanno, MD, PhD<sup>7</sup>; Roisin Colleran, MB, BCh<sup>5,8</sup>; Javier Escaned, MD, PhD<sup>9</sup>; Daniele Giacoppo, MD<sup>10,11</sup>; Vijay Kunadian, MBBS, PhD<sup>12</sup>; Alexandra Lansky, MD<sup>13,14</sup>; Julinda Mehilli, MD<sup>15,16</sup>; Franz-Josef Neumann, MD<sup>17</sup>; Damiano Regazzoli, MD<sup>2</sup>; Jorge Sanz-Sánchez, MD, PhD<sup>1</sup>; William Wijns, MD, PhD<sup>18</sup>; Andreas Baumbach, MD, PhD<sup>13,14</sup>

1. Department of Biomedical Sciences, Humanitas University, Pieve Emanuele-Milan, Italy
2. Humanitas Clinical and Research Center IRCCS, Rozzano-Milan, Italy
3. Cardiac Department, Hospital Universitario de La Princesa, IIS-IP, CIBER-CV, Universidad Autónoma de Madrid, Spain
4. Department of Advanced Biomedical Sciences, University Federico II, Naples, Italy
5. Dublin Cardiovascular Research Institute, Mater Private Hospital, Dublin, Ireland
6. School of Pharmacy and Biomolecular Sciences, Royal College of Surgeons in Ireland, Dublin, Ireland
7. Division of Cardiology, Azienda-Ospedaliero-Universitaria "Policlinico G: Rodolico- San Marco", University of Catania, Italy
8. Department of Cardiovascular Diseases, Deutsches Herzzentrum München, Technische Universität München, Munich, Germany
9. Hospital Clínico San Carlos IDISSC, Complutense University of Madrid, Madrid, Spain
10. Alto Vicentino Hospital, Vicenza, Italy
11. Deutsches Herzzentrum München, Munich, Germany
12. Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University and Freeman Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, United Kingdom.
13. Division of Cardiovascular Medicine, Yale University School of Medicine, New Haven, Connecticut, USA
14. Centre for Cardiovascular Medicine and Devices, William Harvey Research Institute, Queen Mary University of London, and Barts Heart Centre London, United Kingdom
15. Medizinische Klinik und Poliklinik I, Klinikum der Universität München, Ludwig-Maximilians-Universität, Germany
16. German Centre for Cardiovascular Research, partner site Munich Heart Alliance, Munich, Germany
17. Department of Cardiology and Angiology II, University Heart Centre Freiburg-Bad Krozingen, Germany
18. The Lambe Institute for Translational Medicine and Curam, Saolta University Healthcare Group, National University of Ireland Galway, Ireland

**Financial disclosures/Funding:** GGS reports a research grant (to the Institution) from Boston Scientific and speaker/consulting fees from B. Braun, Biosensors, and Boston Scientific. RAB reports lecture fees from B Braun Melsungen AG and Biotronik, and research funding to the institution of employment from Celonova Biosciences. DC declares consulting honoraria from Abbott Vascular, Bayer and Daiichi Sankyo, and speakers' fee from AstraZeneca, Biosensors, Bayer, Boehringer Ingelheim, Daiichi Sankyo, Sanofi Aventis. EB declares speaker's fees from BSCI, Abbott Vascular, GE. VK reports consulting fees/honoraria: Bayer, Amgen, Daiichi Sankyo, Abbot Vascular, Astra Zeneca and major institutional research grant from Astra Zeneca. JM has received lecture fees from Abbott Vascular, Boston Scientific, Biotronik, Edwards Lifesciences, Bristol-Myers Squibb, Medtronic and Terumo; and has received institutional research grants from Edwards Lifesciences, Boston Scientific, and Abbott Vascular. DR reports speaker honoraria from Amgen and Boehringer. AB reports Institutional Research Support from Abbott Vascular and honoraria from Astra Zeneca, Sinomed, Microport, Abbott Vascular, Cardinal Health, KSH.

**\*Address for correspondence:**

Giulio G. Stefanini, MD, PhD, MSc  
Humanitas Clinical and Research Center IRCCS  
Via Manzoni, 56  
20089 Rozzano-Milan, Italy  
email: giulio.stefanini@gmail.com

## Keywords

Prior cardiovascular surgery; Prior PCI; In-stent restenosis; Saphenous vein graft;  
Stent thrombosis; Multidisciplinary Heart Team

## Abbreviations

ACS= acute coronary syndrome

BID= bis in die

BMS= bare-metal stent

CABG= coronary artery bypass grafting

CAD= coronary artery disease

CI= confidence interval

DAPT= dual antiplatelet treatment

DCB= drug-coated balloons

DES= drug-eluting stent

ECG= electrocardiogram

FFR= fractional flow reserve

HR= hazard ratio

ISR= in-stent restenosis

IVUS= intravascular ultrasound

LAD= left anterior descending artery

LIMA= left internal mammary artery anastomosis

MI= myocardial infarction

OCT= optical coherence tomography

OR= odds ratio

PCI= percutaneous coronary interventions

PROSPECT= Prospective Natural History Study of Coronary Atherosclerosis

RR= risk ratio

TLR= target lesion revascularization

## **Abstract**

Myocardial revascularization represents the most frequently performed therapeutic intervention worldwide. Current percutaneous and surgical revascularization techniques provide excellent short- and long-term clinical outcomes. However, despite the technological and procedural advances with the widespread use of drug-eluting stents and arterial bypass grafts in contemporary practice, a considerable proportion of patients require repeat revascularization procedures during long-term follow-up. The need for repeat revascularization has a major impact on patients quality of life and is associated with a significant economic burden. This Consensus Document summarizes the views on the management of myocardial revascularization failure of an expert panel of the European Association of Percutaneous Cardiovascular Interventions (EAPCI). The present document provides a broad and pragmatic overview of the clinical management of myocardial revascularization failure with a focus on the three key underlying mechanisms leading to repeat revascularization: 1) failure of percutaneous coronary interventions, 2) failure of coronary artery bypass grafting and 3) progression of coronary artery disease in native coronary segments previously untreated. The scope of the present position document is to provide a patient-oriented approach for the management of myocardial revascularization failure.

## 1. Introduction

Myocardial revascularization represents the most frequently performed therapeutic intervention worldwide.[1,2] Current revascularization techniques provide excellent clinical outcomes during long-term follow-up.[1,3] Notwithstanding, approximately 20% of patients undergoing myocardial revascularization require a repeat revascularization procedure during the first 5 years of follow-up, with a higher risk after percutaneous coronary interventions (PCI) as compared with coronary artery bypass grafting (CABG).[4–7] The need for repeat revascularization has a significant impact on quality of life and health care resources, and exposes patients to risks intrinsically related to repeat hospitalizations and invasive procedures.[4,8,9] Moreover, patients requiring repeat revascularization are characterized by a high cardiac risk profile, due to comorbidities and anatomical features,[7][10] rendering their clinical management a relevant challenge in daily practice.

This document summarizes the views on the management of myocardial revascularization failure of an expert panel of the European Association of Percutaneous Cardiovascular Interventions (EAPCI). The committee members were proposed by the document chair and co-chair, and approved by the EAPCI Scientific Documents and Initiatives Committee.

This document approaches the management of myocardial revascularization failure from a patient-oriented perspective, based on the underlying mechanisms leading to the clinical need for repeat revascularization: failure of PCI, failure of CABG, and progression of coronary artery disease (CAD) in native coronary segments previously untreated. The latter is not directly related to overt failure of a previous PCI or CABG. However, from a patient perspective, the need for a new revascularization

procedure represents a failure of the initial treatment strategy and should, therefore, be evaluated in the context of revascularization failure.

This document has three key objectives: 1) To outline the different mechanisms underlying myocardial revascularization failure; 2) To detail the specific challenges to short- and long-term success of repeat revascularization procedures; 3) To delineate systematic and informed strategies aiming to increase the safety and efficacy of these procedures.

Copyright EuroIntervention

## 2. Failure of percutaneous coronary interventions

The vast majority of PCI procedures include stent implantation. Stent thrombosis and restenosis are key mechanisms of stent failure requiring repeat revascularization.

### 2.1 Stent thrombosis

#### Early stent thrombosis

Early stent thrombosis is defined as stent thrombosis occurring within the first 30 days after stent implantation and is subclassified into acute (0-24 hours) and subacute (>24 hours–30 days) stent thrombosis.[11] Early stent thrombosis is a relatively infrequent occurrence in contemporary clinical practice (**Table 1**).[12] Most cases are related to mechanical or anatomical factors, in association with a thrombogenic milieu or an acute triggering event (**Table 2**).

#### Late and very late stent thrombosis

Late stent thrombosis is defined as stent thrombosis that occurs between 30 days and one year after stent implantation, while very late stent thrombosis is defined as stent thrombosis that occurs later than year after stent implantation.[13]

In contemporary large-scale drug-eluting stent (DES) trials, with broad inclusion criteria, stent thrombosis rates are low beyond 30 days after stent implantation (**Table 1**).

Risk factors and underlying mechanisms of late and very late stent thrombosis are summarized in **Table 2**.



### Management of patients with stent thrombosis

Most patients with stent thrombosis present with acute myocardial infarction (MI), with or without ST-segment elevation.[14] Accordingly, the principles of management are those recommended in relevant clinical practice guidelines.[15–17] Usually patients with suspected ST should undergo urgent coronary angiography to confirm the diagnosis and treat the underlying cause.

Liberal use of intracoronary imaging [18] – with intravascular ultrasound or optical coherence tomography - is recommended by clinical practice guidelines, in order to detect and modify underlying mechanical factors, and to assess the contribution of concomitant restenosis or neoatherosclerosis to in-stent obstruction.[15]

In case of a completely occluded vessel, flow should be restored initially, and intravascular imaging should be performed afterwards. In addition to intracoronary imaging, radiological stent enhancement is a helpful method to diagnose loss of stent integrity or under-expansion.[19] Although routine thrombus aspiration is not recommended by current guidelines, it might be considered in selected cases of stent thrombosis with a large thrombus burden. Similarly, glycoprotein IIb/IIIa receptor antagonists should be considered in view of the elevated pro-thrombotic milieu. Cangrelor use may be considered in patients not treated with a P2Y12 inhibitor at the time of stent thrombosis.

Identified factors likely to have contributed to stent thrombosis should be corrected (**Figure 1**). Patients with deficits in mechanical stent integrity – such as stent gap, stent fracture or longitudinal deformation – as well as those with residual edge disease or dissection should generally be treated with repeat stenting. Stent crush or collapse is very rare but may be seen in heavily calcified lesions or at ostial locations and

also mandates repeat stenting. Significant stent under-expansion or malapposition should be corrected with non-compliant balloon dilation, including use of balloons with very high rated burst pressure as required. Intravascular lithotripsy may be considered for severe, otherwise non-dilatable stent under-expansion.[20] Following dilation of under-expanded stents, additional stent may be considered, although systematic repeat stenting in such cases should be avoided, especially if there are already multiple stent layers.

Non-mechanical causes of stent thrombosis may predominate in some cases. These include insufficient platelet inhibition due to hyporesponsiveness, non-compliance to antiplatelet therapy, or interruption for unplanned or non-deferrable surgery. In the absence of clear identifiable mechanical causes of stent thrombosis, it may be sufficient to dilate the thrombosed stent to restore blood flow and administer antithrombotic agents (e.g. glycoprotein receptor inhibitors, intravenous P2Y<sub>12</sub> inhibitors). Subsequently, insufficient platelet inhibition must be evaluated. Use of point-of-care phenotypic and genetic testing have been suggested in patients with stent thrombosis without an evident underlying mechanical cause.[21,22] Assessment of dual antiplatelet treatment (DAPT) compliance is of paramount importance, especially within the first 30 days after PCI.[23] Prasugrel and ticagrelor are preferred over clopidogrel after an acute stent thrombosis.[15] Prolonged DAPT therapy beyond 12 months should be considered in patients after a stent thrombosis, weighting their increased thrombotic risk against their bleeding risk.[24–26]

---

### Stent Thrombosis: What To Do

Intracoronary imaging with IVUS and/or OCT to identify factors likely to have contributed to stent thrombosis

PCI with DES in case of deficits in mechanical stent integrity (stent fracture or collapse)

PCI with DES in case of residual edge disease or dissection

High-pressure non-compliant balloon dilation in case of stent under-expansion or malapposition

Assess adherence to antiplatelet therapy

Assess platelet reactivity with point-of care assays in selected cases of acute stent thrombosis without a clearly identified mechanical cause

After PCI for stent thrombosis, dual antiplatelet therapy with aspirin 75-100mg daily and prasugrel 10mg daily or ticagrelor 90mg BID for 12 months

---

### Stent Thrombosis: What NOT To Do

Systematic repeat stenting in cases of stent under-expansion, especially in the presence of multiple stent layers

---

## 2.2 In-stent restenosis

In-stent restenosis (ISR) is a response to vessel wall injury that results in excessive tissue formation (i.e., neointimal hyperplasia or neoatherosclerosis) in stented segment. ISR is an angiographic diagnosis, defined as a diameter stenosis >50% within the stented segment (i.e. the stent and a 5 mm border proximal or distal to the stent). Although DES were highly effective in reducing the risk of ISR compared with bare-metal stent (BMS), ISR remains the most frequent cause of stent failure and the most common indication for target lesion revascularization (TLR). Large-scale clinical trials of patients treated with contemporary DES with broad inclusion criteria report rates of clinical restenosis (i.e. clinically indicated TLR) of <3% at 1-year and 10% at 5 years (**Table 1**). Of note, ISR presents as an acute coronary syndrome in up to 20% of cases.[27]

Clinical and angiographic factors predisposing to ISR are summarized in **Table 2**.

### Management of patients with in-stent restenosis

Treatment of ISR is challenging compared with treatment of *de novo* lesions, owing to relatively high recurrence rates.[28]

As the underlying substrate in ISR often overlaps with that of stent thrombosis, the principles of management are similar. However, while patients with thrombosis usually present with acute MI, patients with ISR may be asymptomatic and should only be treated in the presence of symptoms or objective evidence of ischemia. In stable settings, if revascularization is deemed necessary the strategy should be carefully planned. As is the case with native coronary artery stenoses, when ISR angiographic severity is unclear physiological guidance should be considered. If possible, the original lesion and the initial procedure (e.g., material used, maximum balloon pressures,

challenges encountered, etc.) should be reviewed to identify potential technical issues that may need to be addressed during the repeat intervention. Intracoronary imaging of restenotic lesions, with IVUS or OCT, may provide insights into the mechanisms underlying ISR (**Table 2**), by identifying contributing mechanical factors as well as characterizing the restenotic tissue type. Of note, in addition to intracoronary imaging, radiological stent enhancement is a helpful method to diagnose stent fracture or under-expansion in patients with ISR.[19]

There are a number of technical issues that should be considered in the treatment of patients with ISR. Treatment should generally be focused on the stenosed segment rather than on the full length of the stented segment.[28] To prevent recurrent ISR, it is important to optimize the results of repeat procedures. Careful lesion preparation is required and mechanical issues should be recognized and corrected. Aggressive dilation of the underlying stent might be required especially in under-expanded or collapsed stents, ideally using noncompliant balloons at high pressures (frequently >18 bar). Care should be taken to avoid geographic miss as this may lead to edge-related recurrence. Use of cutting balloons, or more flexible scoring balloons for lesion pre-dilation, reduces slippage of the balloon out of the stent (so called “watermelon seeding”), which may lead to stent edge dissections, with the potential for subsequent “candy wrapper” patterns of stent edge-restenosis. These devices also incise the surface of the neointimal tissue, which theoretically may facilitate the uptake of drug delivery with drug-coated balloons (DCB) angioplasty or repeat DES implantation. Indeed, the ISAR-DESIRE-4 trial, showed improved angiographic outcomes after lesions pre-dilation with a scoring balloon compared with plain balloon angioplasty prior to DCB-angioplasty.[29]

Occlusive ISR constitutes a challenging lesion subset for revascularization. While the use of a contemporary approach to chronic total occlusion recanalization is associated with improved procedural outcomes,[30] long-term results are worse than in de novo chronic total occlusion lesions, largely due to higher TLR rates.[31]

In the case of resistant stent under-expansion, very-high pressure (25 to 35 bar) balloons may be used. Modification of calcific plaques accounting for stent under-expansion can be performed with excimer laser atherectomy [32] or intravascular lithotripsy, the latter also being useful in ISR with calcified neoatherosclerosis.[33] [34] Rotational atherectomy (also termed “rotastenting”) of undilatable under-expanded stents might be considered a second-line strategy and should be undertaken with caution due to the risk of serious complications.[35] Further study of the therapies discussed is required to confirm their potential benefits.

Following lesion preparation, a proportion of patients will require repeat stenting to correct loss of mechanical integrity of the underlying stent (e.g. due to fracture or gap or in rare cases, with demonstrated stent collapse). In the remaining patients, after dilatation and correction of any stent under-expansion, a number of treatment options are available, but there is general consensus that additional treatment beyond mechanical dilatation is required as outcomes after plain balloon angioplasty alone are poor.[36] The two most effective options are DCB-angioplasty or repeat stenting with DES.[37,38] European clinical practice guidelines recommend the use of DES or DCB as first-line therapy in patients with ISR (class I recommendation and level of evidence A for both).[15] Repeat stenting with DES seems to be marginally more effective in terms of angiographic recurrences and need for TLR as compared with DCB, particularly in patients with ISR of DES.[38,39] However, DCB avoids multiple metallic

layers on the vessel wall, which may be of particular concern in patients with recurrent ISR. Accordingly, selection between the two strategies may be considered based on the individual characteristics of the patient and lesion to be treated. For instance, DCB may be preferred over DES in ISR of BMS, multiple metal layers, or large side branches. Conversely, DES may be preferred over DCB in lesions with stent fracture, diffuse ISR extending beyond the stent edges, or in case of significant residual dissection or impaired flow after a balloon-only approach (**Figure 2**). Some operators prefer repeat stenting in the case of ISR at the stent edge though studies suggest that DCB appear to be equally effective for ISR confined to the body of the stent as for those mainly involving its edges.[40,41]

Antiplatelet treatment for patients undergoing PCI for ISR should not differ from that in patients with a de novo lesion. When ISR clinically presents as chronic coronary syndrome, switching antiplatelet therapy is not recommended unless neoatherosclerosis with plaque rupture or erosion is identified by intracoronary imaging.[42]

---

## In-Stent Restenosis: What To Do

Intracoronary imaging with IVUS and/or OCT to detect stent-related mechanical problems leading to ISR

Aggressive pre-dilation of the underlying stent with non-compliant balloons at high pressure, especially in under-expanded or collapsed stents

Lesion preparation with cutting balloons or scoring balloons in order to reduce balloon slippage outside the stent

Very-high pressure balloons, intravascular lithotripsy, excimer laser or rotational atherectomy in case of resistant stent under-expansion

After adequate lesion preparation, PCI with DES or DCB

DES preferred for suboptimal predilation results (residual stenosis >50%, large\* or flow-limiting dissections), diffuse ISR, loss of mechanical integrity, and failed DCB-strategy

DCB preferred for focal ISR, first ISR episode, ISR of BMS, and multiple metal layers

CABG or a conservative strategy instead of a new PCI attempt in patients with recurrent episodes of diffuse ISR, after a Heart Team discussion

After PCI for ACS due to underlying ISR, dual antiplatelet therapy with aspirin 75-100mg daily and prasugrel 10mg daily or ticagrelor 90mg BID for 12 months

---

## In-Stent Restenosis: What NOT To Do

Treatment of the full length of the initial stent instead of focusing on the stenosed segment

Plain balloon angioplasty-only strategy

---

\* Large dissection defined if: longitudinal extension >2mm, lateral extension >60° and involvement of medial or adventitia layers.[43]



### ▪ 2.3. Acute functional failure after PCI

Failing to identify hemodynamically significant coronary stenoses is one of the most common reasons portending to revascularization failure. Complementing coronary angiography with invasive functional assessment has received the highest level of recommendation by current guidelines to evaluate the hemodynamic relevance of intermediate-grade stenosis, when non-invasive evidence of ischemia is not available.[15] Myocardial revascularization aims to eliminate ischemia and is, therefore, expected to normalize findings of invasive functional assessment.

While angiography is considered to be of limited ability to assess the hemodynamic relevance of coronary lesions, the adequacy of acute results after PCI is still mainly assessed based on angiographic visual estimation only. However, early evidence with fractional flow reserve (FFR) suggested that suboptimal FFR after stenting is an independent predictor of adverse clinical outcomes at 6 months.[44] More recently, a prospective observational study including 574 consecutive patients (664 lesions) with FFR pre- and post-PCI evaluated clinical outcomes during a mean follow-up of 31±16 months. Despite adequate angiographic result, 143 lesions (21%) had post-PCI FFR values within the ischemic range (FFR ≤0.80).[45]

A meta-analysis that synthesized evidence of 59 observational (prospective and retrospective) studies evaluating the relationship between post-PCI FFR and clinical outcomes found a normal distribution of post-PCI FFR values, with a mean of 0.90±0.04, and indicated that post-PCI FFR values appear to be related to the risk of repeat revascularization (OR 0.43, 95% CI 0.34-0.56) and major adverse cardiac events (0.71, 95% CI 0.59-0.85) during follow-up.[46] A threshold of final FFR <0.90 has been proposed to define a suboptimal result after stenting.[47]

Several investigations showed that additional interventions may optimise the acute result in patients with suboptimal post-PCI FFR.[45,47,48] A recent prospective small-scale study suggested that intracoronary imaging with optical coherence tomography may reveal potentially treatable causes (i.e., stent underexpansion, incomplete lesion coverage, stent malposition, edge dissection, or tissue protrusion), allowing optimisation of the post-PCI functional result.[47] However, whether additional interventions based on post-PCI functional assessment have a significant impact on clinical outcomes has not been clearly determined.[49]

---

### Acute Functional Failure: What To Do

Repeat invasive functional assessment after stenting when already used to assess the hemodynamic relevance of the treated lesion

Attempt to identify reasons for suboptimal (i.e., FFR <0.90) invasive functional assessment post-PCI, possibly with the use of intracoronary imaging

---

### 3. Failure of coronary artery bypass grafting

Surgical graft failure is frequently observed with increasing time after CABG. Graft failure after use of saphenous vein grafts is as high as 50% at 10 years with vein graft occlusion rates of up to 27% within the first year after CABG.[50–52] Within the first month after surgery, the causes of graft failure are mostly related to the surgical technique and flow-pattern related thrombotic complications, while graft failure thereafter is characterized by neointimal hyperplasia and accelerated progression of CAD.[53–55]

#### 3.1 Acute graft failure (<1 month after surgery)

Acute graft failure can be due to graft dissection, kinking or twisting, anastomotic technical errors, impaired vessel run-off into the native coronary artery, competitive flow from the native coronary artery, or graft thrombosis. In a study of 366 patients with routine post-CABG angiography, 12.2% of the grafts were found to have relevant angiographic defects requiring a minor adjustment of the graft in 2.8%, an anastomosis revision in 3.4%, and intraoperative open-chest PCI in 6.0%.[53] Because of the logistic issues associated with routine direct postoperative angiography, intraoperative transit-time flow measurements and high-frequency epicardial ultrasound have been used to detect causes of graft failure before chest closure and allow the opportunity for revision before myocardial ischemia occurs or progresses.

When clinically relevant, acute graft failure may result in MI with a subsequent risk of mortality. The suspicion of early graft failure should arise in the presence of sudden clinical deterioration as indicated by ECG signs of ischemia, ventricular arrhythmias, biomarker changes, new wall motion abnormalities, or haemodynamic

instability. Due to the low specificity of ECG changes and echocardiographic wall motion abnormalities during the postoperative course and the delay in appearance of biomarker changes, careful assessment of all variables will influence the decision making for angiographic evaluation.[15]

Despite arterial grafting is recommended by current guidelines on myocardial revascularization [15], vein grafts continue to be used in larger numbers than arterial grafts, despite having lower long-term patency rates.[56] Arterial grafts tend to be reserved for the prognostically most important areas of myocardium (e.g., the left internal mammary artery anastomosis [LIMA] to the left anterior descending artery [LAD]). Acute arterial graft failure, therefore, typically has a more severe clinical presentation than vein graft failure, while the latter more often occurs subclinically. An observational study showed that acute graft failure of the LIMA-to-LAD anastomosis warranted reintervention in 80% of patients, while acute vein graft failure was treated conservatively in approximately 50% of patients.[57]

#### Management of acute graft failure

Angiographic assessment is recommended if there is a suspicion of acute graft failure early postoperatively, and is performed in about 1-5% of patients.[58–61] In a recent meta-analysis of 9 studies and 1,104 patients with suspected perioperative MI after CABG, acute graft failure was diagnosed in 62.1% of patients.[62] Incomplete revascularization was the cause of the MI in 6.1% of patients, and 3.5% of patients had a native coronary artery culprit. Remarkably, in 31.6% of patients no cause of perioperative MI could be identified. In this context, it is important to underscore that

haziness at the anastomosis in this acute period may be difficult to interpret and may not be related to the clinical problem.

The treatment strategy for acute graft failure should be made in an ad hoc Heart Team meeting. As summarized in **Figure 3**, a number of parameters should be considered in the decision-making process such as technical reason for acute failure (i.e., problems related with the suture), age and risk profile of the patient, the patient's clinical condition (e.g., haemodynamic status and inotropic support), pre-CABG native vessel CAD and coronary anatomy, extent and timing of ischemia, graft configuration, and extent of myocardium at risk.

In the setting of acute graft failure, emergency PCI may limit the extent of infarction. Current clinical guidelines advocate that PCI should be the preferred strategy in cases of acute graft failure where the anatomy is suitable.[15] In such cases, the target for PCI should be the native vessel or the internal mammary artery (IMA) graft, while an acutely occluded vein graft and any anastomotic site should be avoided, if possible, due to concerns regarding fragility of the new anastomosis as well as the risk of embolisation and perforation. The impact of in-hospital PCI following CABG was investigated in a retrospective study in which patients with acute coronary ischemia requiring PCI after CABG (N=14,323) were compared with those who did not undergo PCI (N=540,664). Post-CABG PCI was associated with an increased risk of unadjusted in-hospital mortality (5.1% vs. 2.7%;  $p < 0.001$ ), higher rates of stroke (2.1% vs. 1.6%;  $p < 0.001$ ), acute kidney injury (16% vs. 12.3%;  $p < 0.001$ ), and a 50% cost increase.[63]

Redo CABG should be preferred when the anatomy is unsuitable for PCI, when an anastomotic error is evident, or when several important grafts are occluded.[15]

Conservative treatment should be considered in cases where diagnosis has been

delayed and viability is expected to be limited. In asymptomatic patients, repeat revascularisation should be considered if the failed graft supplies a large territory of myocardium.

---

### Acute Graft Failure: What To Do

Coronary angiography after CABG in patients with sudden clinical deterioration indicated by:

- symptoms of ischemia and/or abnormal biomarkers suggestive of perioperative MI
- ischemic ECG changes indicating large area of myocardium at risk
- new significant wall motion abnormalities
- haemodynamic instability.

Emergency redo-CABG or PCI decided upon by ad-hoc consultation in the Heart Team, based on the feasibility of revascularization, area at risk, comorbidities, and clinical status

PCI of the native vessel rather than PCI of the graft

Conservative treatment in graft failure cases where diagnosis has been delayed and viability is expected to be limited

In asymptomatic patients, repeat revascularisation if the failed graft supplies a large territory of myocardium

---

### Acute Graft Failure: What NOT To Do

PCI in case of unsuitable anatomy, anastomotic error of the LIMA to LAD or at the Y-anastomosis of a composite arterial graft

---

## 3.2 Late graft failure (>1 month after CABG)

As the time from surgery increases, vein grafts become prone to a process of aggressive and accelerated atherosclerosis. This results in mostly diffuse soft lipid-rich atherosclerotic plaques with extensive necrotic cores with or without intraplaque haemorrhage prone to rupture and downstream embolization.[54][64]

Clinically relevant late graft failure presents mostly in form of stable or unstable angina pectoris.[65–68]

### Management of late graft failure

A number of critical issues should be considered when treating patients with degenerated grafts, including whether to perform redo CABG or PCI, whether to treat native arteries or degenerated grafts, and the risk of distal embolization in case of graft intervention.

PCI is considered treatment of choice in case of late graft failure. Randomized comparisons between redo CABG and PCI, however, are lacking partly due to patients' unwillingness to be allocated to redo CABG.[69] In a subgroup analysis of patients with late graft failure from the AWESOME trial and registry, redo CABG surgery was associated with higher peri-procedural mortality as compared with PCI.[69] Therefore, redo CABG surgery is recommended only in case of extensive native CAD with multiple graft occlusion, particularly in the absence of patent arterial grafts.[15]

PCI of vein grafts is considered a high-risk intervention due to an increased risk of slow/no-reflow related to distal embolization of the friable atheroma, depending on the degree of graft degeneration.[70,71] Embolic protection devices have been proposed to prevent distal embolization.[72,73] A randomized trial performed in early 2000 showed a significant benefit of embolic protection devices in PCI of vein grafts [72] and a similar trend was seen in a subsequent randomized trial that was underpowered due to premature termination [73]. However, a meta-analysis of 52.893 patients enrolled in these randomized trials and in more recent observational studies did not suggest a benefit of routine use of embolic protection devices in PCI of vein grafts.[74]

Several randomized trials have compared DES with BMS in vein graft lesions.[68] In a meta-analysis of randomized evidence, no differences between DES and BMS was

observed in terms of all-cause death (RR 1.06, 95% CI 0.76-1.48), MI (RR 0.81, 95% CI 0.50-1.29), target vessel revascularization (RR 0.73, 95% CI 0.48-1.11) and target lesion revascularization (RR 1.05, 95% CI 0.76-1.43) at longest follow-up.[68] In the ISAR-CABG trial, DES use was associated with a significantly lower risk of target-lesion revascularization during the first year of follow-up (HR 0.49, 95%CI 0.28-0.86) which was offset by a higher risk between 1 and 5 years (HR 2.10, 95% CI 1.37-3.22) as compared to BMS, with a significant interaction between treatment effect and time ( $p_{\text{interaction}} < 0.001$ ) .[66,67]

PCI of vein grafts is associated with a higher risk of adverse events as compared to PCI of native coronary arteries among patients with late graft failure.[75] In a registry of 11,118 veterans, PCI of vein grafts was associated with significantly higher risk of mortality (adjusted hazard ratio 1.30, 95% CI 1.18-1.42), MI (adjusted HR 1.61, 95% CI 1.43-1.82) and repeat revascularization (adjusted HR 1.69, 95% CI 1.50-1.71) as compared to PCI of native arteries during a median follow-up of 3 years.[75]

Although available evidence clearly supports PCI of the native artery in case of late graft failure, anatomical complexities – such as multiple chronic total occlusions of native arteries – might limit the success of such strategy, forcing interventionalists to treat degenerated grafts instead. Despite improvements in recanalization techniques and available dedicated tools, previous CABG surgery remains one of the most important predictors of PCI failure in chronic total occlusions.[76] Therefore, the decision to treat native artery lesions or surgical grafts depends on CAD anatomical complexity and the interventionalists expertise in complex PCI, seeking the most complete revascularization. The decision should be made on an individual patient basis, giving priority to PCI of native arteries.



---

### Late Graft Failure: What To Do

PCI as first choice over redo-CABG for late graft failure

PCI of the native vessel rather than PCI of the graft

PCI strategy based on operator experience in complex PCI

Distal protection devices for PCI of vein graft lesions with diffused degeneration

IMA for redo-CABG in patients in whom the IMA was not used previously

Redo-CABG in patients without a patent IMA graft to the LAD, after checking its patency

Redo-CABG in case of extensive native CAD, anatomically unsuitable for PCI, in absence of patent grafts (especially arterial)

---

### Late Graft Failure: What NOT To Do

Routine use of embolic protection devices for PCI of vein grafts

Plain balloon-only for PCI of the graft

---

#### 4. Repeat revascularization due to progression of CAD

CAD progression in native coronary segments previously untreated is the primary cause of repeat procedures after myocardial revascularization.

##### Native CAD progression after PCI

Disease progression is responsible for a relevant proportion of repeat revascularization procedures after PCI,[6] although the incidence varies based on the clinical and anatomic characteristics of the population studied. The Prospective Natural History Study of Coronary Atherosclerosis (PROSPECT) study studied the relative contribution of events related to the initially treated lesion (culprit lesion) and events related to CAD progression in non-culprit sites among 697 patients with ACS undergoing PCI.[77] The cumulative rate of major adverse cardiac events – a composite of cardiac death, arrest, MI, and hospitalization for angina – was 20.4% at 3 years, with 12.9% events related to the culprit lesion and 11.6% events due to CAD progression at non-culprit sites. Overall, 65% of all events occurred within 1 year after PCI, with a relatively equal distribution between events related to the culprit lesion and those related to CAD progression. The overall repeat revascularization rate was 17.1% at 3 years, with an equal contribution of events related to the culprit lesion and those related to CAD progression.

Predictors of CAD progression in previously untreated native coronary segments include clinical and angiographic factors that are largely overlapping with predictors of PCI and CABG failure, such as age, diabetes mellitus, complex coronary anatomy, extent of CAD, small vessel CAD, and previous PCI of vein grafts or ostial lesions.[6][78]

### Native CAD progression after CABG

Current recommendations for CABG inherently select patients at higher risk for native CAD progression. These include patients with multi-vessel CAD, high anatomical complexity and extent of CAD, and coexistence of multiple comorbidities including diabetes mellitus, reduced left ventricular ejection fraction, and chronic kidney disease. Historical evidence indicates that accelerated CAD progression occurs up to 10-fold more frequently in non-obstructive atherosclerotic lesions in bypassed coronary arteries compared with similar lesions in non-bypassed vessels at 3 years after CABG.[79] In another study, the risk of CAD progression was twice as high in arteries with patent grafts as compared to those with closed grafts, with the majority of grafted arteries with CAD progression being completely occluded. A more recent analysis of contemporary surgical techniques showed similar results, with development of a new chronic total occlusion in a native coronary artery in >40% of patients within 1 year after CABG, strongly predicted by a severe (>90%) proximal stenosis in the same vessel.[80]

### General principles for management of CAD progression

In case of CAD progression in previously untreated native coronary segments following revascularization, treatment recommendations should be based on symptoms and evidence of myocardial ischemia. In this context, optimal medical therapy plays a pivotal role not only to reduce the risk of CAD progression but also for an initial management of patients with evidence of CAD progression. We refer to relevant clinical practice guidelines for a comprehensive assessment on recommendations for optimal medical management, which represents the cornerstone for prevention and treatment

of CAD progression.[15,81,82] The interventional management of CAD progression differs according to the initial revascularization modality.

#### Management of CAD progression after PCI

In contemporary large-scale PCI trials, up to one-third of patients enrolled were previously treated with PCI.[83–85] Percutaneous treatment of CAD progression after a previous PCI is generally reasonable. A surgical revascularization strategy may be appropriate in case of CAD progression involving proximal segments of major coronary arteries or multivessel disease involving the left main or proximal LAD. A large registry that evaluated outcomes of patients with previous PCI undergoing CABG showed that early mortality and adverse ischemic events did not significantly increase in patients with single or multiple previous PCI procedures.[86] Therefore, a strategy based on clinical and anatomical factors similar to patients with a first diagnosis of CAD is recommended in patients with CAD progression after PCI (**Figure 4**).

#### Management of CAD progression after CABG

Repeat revascularization procedures after CABG are typically performed in older patients with more comorbidities and more complex coronary anatomy as compared to patients with a first diagnosis of CAD. Furthermore, in these patients arterial conduits tend to be less frequently available, having already been used.[87] Therefore, re-do CABG is associated with increased procedural risks and worse clinical outcomes compared with a first CABG. Recent evidence indicates a trend towards a decreased risk of adverse events in patients treated with PCI coupled with an increase in PCI use in this setting.[70][88] In view of the paucity of available comparative effectiveness evidence,

in these patients the selection of the repeat revascularization strategy should be based on the assessment of clinical and anatomical risk profiles on an individual patient basis in discussion within the Heart Team (**Figure 3, Figure 4**).[89][69]

---

### CAD Progression: What To Do

Repeat revascularization in patients with evidence of CAD progression and with a large area of ischemia or severe symptoms despite medical therapy

Base the selection of the repeat revascularization strategy on the assessment of clinical and anatomical risk profiles on an individual basis in the context of the Heart Team

If considered safe, PCI with DES as first choice over CABG

IMA for redo-CABG in patients in whom the IMA was not used previously

Redo-CABG in patients without a patent IMA graft to the LAD

---

### CAD Progression: What NOT To Do

Routine Invasive angiography tests in asymptomatic patients with prior revascularization

Routine ad-hoc PCI in patients with progression of CAD after CABG

---

### Conclusions

Current percutaneous and surgical revascularization techniques are associated with excellent procedural and long-term clinical outcomes. However, a considerable proportion of patients require repeat revascularization procedures during long-term follow-up due to failure of the initial revascularization – either PCI or CABG – or progression of disease in previously untreated coronary segments. This document provides an evidence-based guidance for the management of myocardial revascularization failure based on the underlying mechanism, the timing and the clinical and angiographic characteristics of individual patients.

*Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal*

## 5. References

References available in the online supplementary material.

### Figure legends

#### Figure 1. Algorithm for the management of stent thrombosis.

PCI=percutaneous coronary interventions, DES=drug-eluting stent. \*Avoiding stent implantation should be considered in cases with severe under-expansion or malapposition without further underlying mechanisms. In patients with stent thrombosis due to severe neointimal hyperplasia or neoatherosclerosis PCI with DEB might be considered. Images were kindly provided by Drs Nicolas Amabile, Fernando Alfonso and Gennaro Sardella.

#### Figure 2. Algorithm for the management of in-stent restenosis

BMS=bare-metal stent; DCB=drug-coated balloon. DES=drug-eluting stent; ISR=in-stent restenosis; NC=non-compliant; PCI=percutaneous coronary intervention. \*In patients with edge dissection or acute recoil after lesion predilation, PCI with DES should be considered. Images were kindly provided by Drs Nicolas Amabile, Fernando Alfonso and Gennaro Sardella.

#### Figure 3. Algorithm for the management of patients with surgical graft failure.

CTO= chronic total occlusion; DAPT= dual antiplatelet therapy; LAD= left anterior descending artery; LIMA= left internal mammary artery; PCI= percutaneous coronary intervention; SGF=surgical graft failure

**Figure 4. Factors that may guide revascularization strategy for CAD progression.**

CAD= coronary artery disease; COPD= chronic obstructive pulmonary disease; DAPT= dual antiplatelet therapy; STS = Society of Thoracic Surgery Risk Score

Copyright EuroIntervention

**Table 1. Rates of stent thrombosis and TLR in selected contemporary large-scale all-comer clinical trials[90,91,100–107,92–99]**

	RESOLUTE All comers		DUTCH PEERS		CENTURY II		BIO SCIENCE		BIO-RESORT		SORT OUT VII		BIONICS		BIONYX		SORT OUT VIII		
	Resolute	Xience	Resolute	Promus	Ultimaster	Xience	Orsiro	Xience	Synergy	Resolute	Orsiro	Orsiro	Nobiro	BioNIR	Resolute	Resolute	Orsiro	BioMatrix	Synergy
<b>Definite ST</b>																			
30 days							0.3%	0.2%										0.4%	0.7%
1 year	1.2%	0.3%	0.3%	0.6%	0.9%*	0.9%*	0.9%	0.4%	0.3%	0.2%	0.3%	0.4%	1.2%	0.4%	0.5%	0.1%	0.5%	0.9%	0.7%
2 years			0.8%	0.9%			1.0%	0.7%	0.6%	0.5%	0.4%	0.8%	1.4%						
3 years									0.7%	0.5%	0.7%								
5 years	1.6%	0.8%	1.1%	1.1%	1.3%	1.1%	1.5%	1.5%											
<b>Definite or Probable ST</b>																			
30 days	1.1%	0.5%					1.8%	2.2%											
1 year	1.7%	0.7%	0.5%	0.8%			2.8%	3.4%	0.4%	0.5%	0.4%	0.9%	1.6%	0.4%	0.6%	0.1%	0.7%	1.4%	1.1%
2 years	1.9%	0.9%	1.1%	1.1%			3.7%	4.7%	0.9%	0.7%	0.6%	1.3%	1.8%						
3 years									1.0%	0.8%	1.0%								
5 years	2.4%	1.7%	1.5%	1.3%	1.3%	1.3%	5.8%	7.2%											
<b>TLR</b>																			
30 days																			
1 year			2.2%	2.2%	2.2%*	1.6%*	4.0%	3.1%	1.4%	1.4%	1.5%	2.0%	2.9%	3.2%	2.3%	2.5%	1.9%	2.5%	2.3%
2 years	5.7%	5.1%	3.8%	3.5%			5.5%	4.8%	2.3%	2.9%	2.1%	3.6%	4.5%						
3 years									3.1%	3.6%	2.8%								
5 years	10.2%	8.9%			9.4%	8.2%	10.3%	10.0%											

ST=stent thrombosis, TLR=target lesion revascularization. \*9 months follow-up.



**Table 2. Risk factors and underlying mechanism of stent thrombosis and in-stent restenosis**

	Stent Thrombosis[12,14,18,108–113]			In-Stent Restenosis[111,114–117]
	Early	Late	Very late	
<b>Risk factors</b>				
Patient-related	<ul style="list-style-type: none"> <li>- Acute clinical presentation</li> <li>- Poor response to antiplatelet treatment</li> <li>- High on-treatment platelet reactivity</li> <li>- Current smoking</li> <li>- Genetic variants*</li> <li>- Diabetes mellitus</li> <li>- LVEF &lt;40%</li> </ul>	<ul style="list-style-type: none"> <li>- Current smoking</li> <li>- Multivessel disease</li> <li>- Younger age</li> <li>- LVEF &lt;40%</li> <li>- eGFR&lt;30 ml/(min·m<sup>2</sup>)</li> </ul>	<ul style="list-style-type: none"> <li>- Current smoking</li> <li>- Multivessel disease</li> <li>- Younger age</li> </ul>	<ul style="list-style-type: none"> <li>- Diabetes mellitus</li> <li>- Prior bypass surgery</li> </ul>
Lesion-related	<ul style="list-style-type: none"> <li>- LMCA or LAD lesion</li> <li>- Residual dissection</li> <li>- TIMI flow grade &lt;3</li> <li>- Bifurcation lesion</li> <li>- Type C lesions</li> <li>- Severe calcified lesions</li> </ul>	<ul style="list-style-type: none"> <li>- LAD lesion</li> <li>- Bypass graft lesion</li> <li>- Presence of thrombus</li> <li>- Bifurcation lesion</li> <li>- Severe calcified lesions</li> </ul>	<ul style="list-style-type: none"> <li>- LAD lesion</li> <li>- Bypass graft lesion</li> <li>- Presence of thrombus</li> </ul>	<ul style="list-style-type: none"> <li>- Small vessel size</li> <li>- Complex morphology</li> <li>- Previous diffuse ISR</li> <li>- Bifurcation lesion</li> </ul>
Stent-related	<ul style="list-style-type: none"> <li>- Under-sizing</li> <li>- Overlapping stents</li> </ul>	<ul style="list-style-type: none"> <li>- Long stent length</li> <li>- Overlapping stents</li> </ul>	<ul style="list-style-type: none"> <li>- Long stent length</li> <li>- Overlapping stents</li> </ul>	<ul style="list-style-type: none"> <li>- Long stent length</li> </ul>
<b>Underlying mechanisms</b>	<ul style="list-style-type: none"> <li>- Uncovered struts</li> <li>- Stent under-expansion</li> <li>- Malapposition</li> </ul>	<ul style="list-style-type: none"> <li>- Uncovered struts</li> <li>- Malapposition</li> </ul>	<ul style="list-style-type: none"> <li>- Neoatherosclerosis</li> <li>- Uncovered struts</li> <li>- Malapposition</li> </ul>	<ul style="list-style-type: none"> <li>- Neointimal hyperplasia</li> <li>- Neoatherosclerosis</li> <li>- Stent under-expansion</li> <li>- Loss of mechanical integrity</li> </ul>

eGFR= estimated glomerular filtration rate, ISR=in-stent restenosis, LAD=left anterior descending artery, LMCA= left main coronary artery, LVEF= left ventricular ejection fraction, TIMI=Thrombolysis in Myo

Figure 1




Figure 2

Copyright EuroIntervention




Figure 3





Figure 4



## Supplementary Material

### References

1. Head SJ, Milojevic M, Taggart DP, Puskas JD. Current Practice of State-of-the-Art Surgical Coronary Revascularization. *Circulation*. 2017;136:1331–45.
2. Stefanini GG, Holmes DR. Drug-Eluting Coronary-Artery Stents. *N Engl J Med*. 2013;368:254–65.
3. Byrne RA, Serruys PW, Baumbach A, Escaned J, Fajadet J, James S, Joner M, Oktay S, Jüni P, Kastrati A, Sianos G, Stefanini GG, Wijns W, Windecker S. Report of a European Society of Cardiology-European Association of Percutaneous Cardiovascular Interventions task force on the evaluation of coronary stents in Europe: executive summary. *Eur Heart J*. 2015;36:2608–20.
4. Parasca CA, Head SJ, Milojevic M, Mack MJ, Serruys PW, Morice M-C, Mohr FW, Feldman TE, Colombo A, Dawkins KD, Holmes DR, Kappetein PA. Incidence, Characteristics, Predictors, and Outcomes of Repeat Revascularization After Percutaneous Coronary Intervention and Coronary Artery Bypass Grafting. *JACC Cardiovasc Interv*. 2016;9:2493–507.
5. Fosbøl EL, Zhao Y, Shahian DM, Grover FL, Edwards FH, Peterson ED. Repeat Coronary Revascularization After Coronary Artery Bypass Surgery in Older Adults. *Circulation*. 2013;127:1656–63.
6. Taniwaki M, Stefanini GG, Silber S, Richardt G, Vranckx P, Serruys PW, Buszman PE, Kelbaek H, Windecker S. 4-Year Clinical Outcomes and Predictors of Repeat Revascularization in Patients Treated With New-Generation Drug-Eluting Stents. *J Am Coll Cardiol*. 2014;63:1617–25.
7. Giustino G, Serruys PW, Mehran R, Maehara A, Puskas JD, Simonton CA, Lembo NJ, Kandzari DE, Morice M, Taggart DP, Gershlick AH, Kron IL, Liu Y, Zhang Z, McAndrew T, Dressler O, Ben-yehuda O, Pocock SJ, Kappetein AP, Stone GW. Mortality After Repeat Revascularization Following PCI or Coronary Artery Bypass Grafting for Left Main Disease: The EXCEL trial. *JACC Cardiovasc Interv*. 2020;13(3):375-387.
8. Arnold S V, Magnuson EA, Wang K, Serruys PW, Kappetein AP, Mohr FW. Do Differences in Repeat Revascularization Explain the Antianginal Benefits of Bypass Surgery Versus Percutaneous Coronary

Intervention? Implications for Future Treatment Comparisons. *Circ Cardiovas Qual Outcomes*. 2012 May; 5(3):267-75.

9. Palmerini T, Riva D Della, Biondi-zoccai G, Leon MB, Serruys PW, Smits PC, Ben-yehuda O, Bruno AG, Jenkins P, Stone GW. Mortality Following Nonemergent, Uncomplicated Target Lesion Revascularization After Percutaneous Coronary Intervention. *JACC Cardiovasc Interv*. 2018;11(9):892-902.
10. Escaned J. Secondary coronary revascularisation: an emerging issue. *EuroIntervention*. 2009;5 Suppl D:D6–13.
11. Garcia-Garcia HM, McFadden EP, Farb A, Mehran R, Stone GW, Spertus J, Onuma Y, Morel M, van Es G-A, Zuckerman B, Fearon WF, Taggart D, Kappetein A-P, Krucoff MW, Vranckx P, Windecker S, Cutlip D, Serruys PW, Academic Research Consortium. Standardized End Point Definitions for Coronary Intervention Trials. *Eur Heart J*. 2018;39:2192–207.
12. Byrne RA, Joner M, Kastrati A. Stent thrombosis and restenosis: what have we learned and where are we going? The Andreas Grüntzig Lecture ESC 2014. *Eur Heart J*. 2015;36:3320–31.
13. Garcia-Garcia HM, McFadden EP, Farb A, Mehran R, Stone GW, Spertus J, Onuma Y, Morel M, van Es G-A, Zuckerman B, Fearon WF, Taggart D, Kappetein A-P, Krucoff MW, Vranckx P, Windecker S, Cutlip D, Serruys PW, Academic Research Consortium. Standardized End Point Definitions for Coronary Intervention Trials. *Eur Heart J*. 2018;39:2192–207.
14. Adriaenssens T, Joner M, Godschalk TC, Malik N, Alfonso F, Xhepa E, De Cock D, Komukai K, Tada T, Cuesta J, Sirbu V, Feldman LJ, Neumann F-J, Goodall AH, Heestermans T, Buyschaert I, Hlinomaz O, Belmans A, Desmet W, ten Berg JM, Gershlick AH, Massberg S, Kastrati A, Guagliumi G, Byrne RA. Optical Coherence Tomography Findings in Patients With Coronary Stent Thrombosis. *Circulation*. 2017;136:1007–21.
15. Neumann F-J, Sousa-Uva M, Ahlsson A, Alfonso F, Banning AP, Benedetto U, Byrne RA, Collet J-P, Falk V, Head SJ, Jüni P, Kastrati A, Koller A, Kristensen SD, Niebauer J, Richter DJ, Seferović PM, Sibbing D, Stefanini GG, Windecker S, Yadav R, Zembala MO. 2018 ESC/EACTS Guidelines on myocardial revascularization. *EuroIntervention*. 2019;14:1435–534.



16. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, Caforio ALP, Crea F, Goudevenos JA, Halvorsen S, Hindricks G, Kastrati A, Lenzen MJ, Prescott E, Roffi M, Valgimigli M, Varenhorst C, Vranckx P, Widimský P, ESC Scientific Document Group. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*. 2018;39:119–77.
17. Roffi M, Valgimigli M, Bax JJ, Borger MA, Gencer B, Germany UL, Germany JM, Uk RFS, Windecker S, ESC Scientific Document Group. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2016;37(3):267–315.
18. Souteyrand G, Amabile N, Mangin L, Chabin X, Meneveau N, Cayla G, Vanzetto G, Barnay P, Trouillet C, Rioufol G, Rangé G, Teiger E, Delaunay R, Dubreuil O, Lhermusier T, Mulliez A, Levesque S, Belle L, Caussin C, Motreff P. Mechanisms of stent thrombosis analysed by optical coherence tomography: insights from the national PESTO French registry. *Eur Heart J*. 2016;37:1208–16.
19. Kim MS, Eng MH, Hudson PA, Garcia JA, Wink O, Messenger JC, Carroll JD. Coronary stent fracture: Clinical use of image enhancement. *JACC Cardiovasc Imaging*. 2010;3:446–7.
20. Salazar C, Escaned J, Tirado G, Gonzalo N. Undilatable Calcific Coronary Stenosis Causing Stent Underexpansion and Late Stent Thrombosis: A Complex Scenario Successfully Managed With Intravascular Lithotripsy. *JACC Cardiovasc Interv*. 2019;12:1510–2.
21. Rajendran S, Parikh D, Shugman I, Bch MB, French JK, Chb MB, Juergens CP. High on Treatment Platelet Reactivity and Stent Thrombosis. *Heart, Lung Circ*. 2011;20:525–31.
22. Mega JL, Simon T, Collet JP, Anderson JL, Antman EM, Bliden K, Cannon CP, Danchin N, Giusti B, Gurbel P, Horne BD, Hulot JS, Kastrati A, Montalescot G, Neumann FJ, Shen L, Sibbing D, Steg PG, Trenk D, Wiviott SD, Sabatine MS. Reduced-function CYP2C19 genotype and risk of adverse clinical outcomes among patients treated with clopidogrel predominantly for PCI: A meta-analysis. *JAMA*. 2010;304:1821–30.

23. Dangas GD, Claessen BE, Mehran R, Xu K, Stone GW. Stent thrombosis after primary angioplasty for STEMI in relation to non-adherence to dual antiplatelet therapy over time: results of the HORIZONS-AMI trial. *EuroIntervention*. 2013;8:1033–9.
24. Bonaca MP, Bhatt DL, Cohen M, Steg PG, Storey RF, Jensen EC, Magnani G, Bansilal S, Fish MP, Im K, Bengtsson O, Ophuis TO, Budaj A, Theroux P, Ruda M, Hamm C, Goto S, Spinar J, Nicolau JC, Kiss RG, Murphy SA, Wiviott SD, Held P, Braunwald E, Sabatine MS. Long-Term Use of Ticagrelor in Patients with Prior Myocardial Infarction. *N Engl J Med*. 2015;372:1791–800.
25. Mauri L, Kereiakes DJ, Yeh RW, Driscoll-Shempp P, Cutlip DE, Steg PG, Normand S-LT, Braunwald E, Wiviott SD, Cohen DJ, Holmes DR, Krucoff MW, Hermiller J, Dauerman HL, Simon DI, Kandzari DE, Garratt KN, Lee DP, Pow TK, Ver Lee P, Rinaldi MJ, Massaro JM. Twelve or 30 months of dual antiplatelet therapy after drug-eluting stents. *N Engl J Med*. 2014;371:2155–66.
26. Stefanescu Schmidt AC, Kereiakes DJ, Cutlip DE, Yeh RW, D'Agostino RB, Massaro JM, Hsieh W-H, Mauri L. Myocardial Infarction Risk After Discontinuation of Thienopyridine Therapy in the Randomized DAPT Study (Dual Antiplatelet Therapy). *Circulation*. 2017;135:1720–32.
27. Stone GW, Ellis SG, Colombo A, Dawkins KD, Grube E, Cutlip DE, Friedman M, Baim DS, Koglin J. Offsetting Impact of Thrombosis and Restenosis on the Occurrence of Death and Myocardial Infarction After Paclitaxel-Eluting and Bare Metal Stent Implantation 2007:2842–7.
28. Alfonso F, Byrne RA, Rivero F, Kastrati A. Current Treatment of In-Stent Restenosis. *J Am Coll Cardiol*. 2014;63:2659–73.
29. Kufner S, Joner M, Schneider S, Tölg R, Zrenner B, Repp J, Starkmann A, Xhepa E, Ibrahim T, Cassese S, Fusaro M, Ott I, Hengstenberg C, Schunkert H, Abdel-Wahab M, Laugwitz K-L, Kastrati A, Byrne RA. Neointimal Modification With Scoring Balloon and Efficacy of Drug-Coated Balloon Therapy in Patients With Restenosis in Drug-Eluting Coronary Stents. *JACC Cardiovasc Interv*. 2017;10:1332–40.
30. De La Torre Hernandez JM, Rumoroso JR, Subinas A, Gonzalo N, Ojeda S, Pan M, Yuste VM, Suárez A, Hernández F, Teruel L, Moreu J, Cubero JM, Cascón JD, Vinhas H, Lozano I, Moreiras JM, De Prado AP, Goicolea J, Escaned J. Percutaneous intervention in chronic total coronary occlusions caused by in-stent

- restenosis: Procedural results and long-term clinical outcomes in the TORO (Spanish registry of chronic Total occlusion secondary to an occlusive in-stent RestenOsis) multic. *EuroIntervention*. 2017;13:e219–26.
31. Azzalini L, Dautov R, Ojeda S, Benincasa S, Bellini B, Giannini F, Chavarría J, Pan M, Carlino M, Colombo A, Rinfret S. Procedural and Long-Term Outcomes of Percutaneous Coronary Intervention for In-Stent Chronic Total Occlusion. *JACC Cardiovasc Interv*. 2017;10:892–902.
32. Yin D, Maehara A, Mezzafonte S, Moses JW, Mintz GS, Shlofmitz RA. Excimer Laser Angioplasty-Facilitated Fracturing of Napkin-Ring Peri-Stent Calcium in a Chronically Underexpanded Stent: Documentation by Optical Coherence Tomography. *JACC Cardiovasc Interv*. 2015;8:e137–9.
33. Alfonso F, Bastante T, Antuña P, de la Cuerda F, Cuesta J, García-Guimaraes M, Rivero F. Coronary Lithoplasty for the Treatment of Undilatable Calcified De Novo and In-Stent Restenosis Lesions. *JACC Cardiovasc Interv*. 2019;12:497–9.
34. Salazar C, Escaned J, Tirado G, Gonzalo N. Recurrent restenosis caused by severe calcific neoatherosclerosis treated with intravascular lithotripsy. *EuroIntervention*. 2019.
35. Akin I, Pohlmann S, Nienaber CA, Ince H. A different way of coronary lesion preparation: stentablation and rotastenting. *Clin Med Insights Cardiol*. 2012;6:53–6.
36. Byrne RA, Neumann F-J, Mehilli J, Piniček S, Wolff B, Tiroch K, Schulz S, Fusaro M, Ott I, Ibrahim T, Hausleiter J, Valina C, Pache J, Laugwitz K-L, Massberg S, Kastrati A. Paclitaxel-eluting balloons, paclitaxel-eluting stents, and balloon angioplasty in patients with restenosis after implantation of a drug-eluting stent (ISAR-DESIRE 3): a randomised, open-label trial. *Lancet*. 2013;381:461–7.
37. Siontis GCM, Stefanini GG, Mavridis D, Siontis KC, Alfonso F, Pérez-Vizcayno MJ, Byrne RA, Kastrati A, Meier B, Salanti G, Juni P, Windecker S. Percutaneous coronary interventional strategies for treatment of in-stent restenosis: a network meta-analysis. *Lancet*. 2015;386:655–64.
38. Giacoppo D, Gargiulo G, Aruta P, Capranzano P, Tamburino C, Capodanno D. Treatment strategies for coronary in-stent restenosis: systematic review and hierarchical Bayesian network meta-analysis of 24 randomised trials and 4880 patients. *BMJ*. 2015;351:h5392.

39. Giacoppo D, Alfonso F, Xu B, Claessen BEPM, Adriaenssens T, Jensen C, Pérez-Vizcayno MJ, Kang D-Y, Degenhardt R, Pleva L, Baan J, Cuesta J, Park D-W, Schunkert H, Colleran R, Kukla P, Jiménez-Quevedo P, Unverdorben M, Gao R, Naber CK, Park S-J, Henriques JPS, Kastrati A, Byrne RA. Paclitaxel-coated balloon angioplasty vs. drug-eluting stenting for the treatment of coronary in-stent restenosis: a comprehensive, collaborative, individual patient data meta-analysis of 10 randomized clinical trials (DAEDALUS study). *Eur Heart J*. 2019.
40. Alfonso F, Pérez-Vizcayno MJ, García del Blanco B, Otaegui I, Masotti M, Zueco J, Velázquez M, Sanchís J, García-Touchard A, Lázaro-García R, Moreu J, Bethencourt A, Cuesta J, Rivero F, Cárdenas A, Gonzalo N, Jiménez-Quevedo P, Fernández C. Long-Term Results of Everolimus-Eluting Stents Versus Drug-Eluting Balloons in Patients With Bare-Metal In-Stent Restenosis. *JACC Cardiovasc Interv*. 2016;9:1246–55.
41. Alfonso F, Pérez-Vizcayno MJ, Cárdenas A, García del Blanco B, García-Touchard A, López-Minguéz JR, Benedicto A, Masotti M, Zueco J, Iñiguez A, Velázquez M, Moreno R, Mainar V, Domínguez A, Pomar F, Melgares R, Rivero F, Jiménez-Quevedo P, Gonzalo N, Fernández C, Macaya C. A Prospective Randomized Trial of Drug-Eluting Balloons Versus Everolimus-Eluting Stents in Patients With In-Stent Restenosis of Drug-Eluting Stents. *J Am Coll Cardiol*. 2015;66:23–33.
42. Johnson TW, Räber L, di Mario C, Bourantas C, Jia H, Mattesini A, Gonzalo N, de la Torre Hernandez JM, Prati F, Koskinas K, Joner M, Radu MD, Erlinge D, Regar E, Kunadian V, Maehara A, Byrne RA, Capodanno D, Akasaka T, Wijns W, Mintz GS, Guagliumi G. Clinical use of intracoronary imaging. Part 2: acute coronary syndromes, ambiguous coronary angiography findings, and guiding interventional decision-making: an expert consensus document of the European Association of Percutaneous Cardiovascular Interventions. *Eur Heart J*. 2019;40:2566–84.
43. Räber L, Mintz GS, Koskinas KC, Johnson TW, Holm NR, Onuma Y, Radu MD, Joner M, Yu B, Jia H, Meneveau N, De La Torre Hernandez JM, Escaned J, Hill J, Prati F, Colombo A, Di Mario C, Regar E, Capodanno D, Wijns W, Byrne RA, Guagliumi G, Alfonso F, Bhindi R, Ali Z, Carter R. Clinical use of intracoronary imaging. Part 1: Guidance and optimization of coronary interventions. An expert consensus

document of the European Association of Percutaneous Cardiovascular Interventions. Eur Heart J. 2018;39:3281–300.

44. Pijls NHJ, Klauss V, Siebert U, Powers E, Takazawa K, Fearon WF, Escaned J, Tsurumi Y, Akasaka T, Samady H, De Bruyne B. Coronary Pressure Measurement After Stenting Predicts Adverse Events at Follow-Up: A Multicenter Registry. Circulation. 2002;105(25):2950–4.

45. Agarwal SK, Kasula S, Hacioglu Y, Ahmed Z, Uretsky BF, Hakeem A. Utilizing Post-Intervention Fractional Flow Reserve to Optimize Acute Results and the Relationship to Long-Term Outcomes. JACC Cardiovasc Interv. 2016;9:1022–31.

46. Rimac G, Fearon WF, Bruyne B De, Ikeno F, Matsuo H, Piroth Z, Costerousse O, Bertrand OF. Clinical Value of Post-Percutaneous Coronary Intervention Fractional Flow Reserve Value: A Systematic Review and Meta-Analysis. Am Heart J. 2017;183:1-9.

47. Wolfrum M, Maria GL De, Benenati S, Langrish J, Lucking AJ, Channon KM, Kharbanda RK, Banning AP. What are the causes of a suboptimal FFR after coronary stent deployment? Insights from a consecutive series using OCT imaging. Eurointervention. 2018;14(12):e1324-e1331.

48. Azzalini L, Poletti E, Demir OM, Ancona MB, Mangieri A, Giannini F, Carlino M, Chieffo A, Montorfano M, Colombo A, Latib A. Impact of Post-Percutaneous Coronary Intervention Fractional Flow Reserve Measurement on Procedural Management and Clinical Outcomes: The REPEAT-FFR Study. J Invasive Cardiol. 2019;31(8):229–34.

49. Crea F, Bairey Merz CN, Beltrame JF, Berry C, Camici PG, Kaski JC, Ong P, Pepine CJ, Sechtem U, Shimokawa H. Mechanisms and diagnostic evaluation of persistent or recurrent angina following percutaneous coronary revascularization. Eur Heart J. 2019;40:2455–62.

50. Mehta RH, Ferguson TB, Lopes RD, Hafley GE, Mack MJ, Kouchoukos NT, Gibson CM, Harrington RA, Califf RM, Peterson ED, Alexander JH. Saphenous Vein Grafts With Multiple Versus Single Distal Targets in Patients Undergoing Coronary Artery Bypass Surgery. Circulation. 2011;124:280–8.

51. Lamy A, Eikelboom J, Sheth T, Connolly S, Bosch J, Fox KAA, Zhu J, Lonn E, Dagenais G, Widimsky P, Branch KRH, Bhatt DL, Zheng Z, Straka Z, Dagenais F, Kong Y, Marsden T, Lee SF, Copland I, Yusuf S. Rivaroxaban, Aspirin, or Both to Prevent Early Coronary Bypass Graft Occlusion: The COMPASS-CABG Study. *J Am Coll Cardiol.* 2019;73:121–30.
52. Goldman S, Zadina K, Moritz T, Ovitt T, Sethi G, Copeland JG, Thottapurathu L, Krasnicka B, Ellis N, Anderson RJ, Henderson W. Long-term patency of saphenous vein and left internal mammary artery grafts after coronary artery bypass surgery: Results from a Department of Veterans Affairs Cooperative Study. *J Am Coll Cardiol.* 2004;44:2149–56.
53. Zhao DX, Leacche M, Balaguer JM, Boudoulas KD, Damp JA, Greelish JP, Byrne JG, Writing Group of the Cardiac Surgery, Cardiac Anesthesiology, and Interventional Cardiology Groups at the Vanderbilt Heart and Vascular Institute, Ahmad RM, Ball SK, Cleator JH, Deegan RJ, Eagle SS, Fong PP, Fredi JL, Hoff SJ, Jennings HS, McPherson JA, Piana RN, Pretorius M, Robbins MA, Slosky DA, Thompson A. Routine Intraoperative Completion Angiography After Coronary Artery Bypass Grafting and 1-Stop Hybrid Revascularization. *J Am Coll Cardiol.* 2009;53:232–41.
54. Yahagi K, Kolodgie FD, Otsuka F, Finn AV, Davis HR, Joner M, Virmani R. Pathophysiology of native coronary, vein graft, and in-stent atherosclerosis. *Nat Rev Cardiol.* 2016;13:79–98.
55. Harskamp RE, Lopes RD, Baisden CE, de Winter RJ, Alexander JH. Saphenous Vein Graft Failure After Coronary Artery Bypass Surgery. *Ann Surg.* 2013;257:824–33.
56. Head SJ, Milojevic M, Taggart DP, Puskas JD. Current Practice of State-of-the-Art Surgical Coronary Revascularization. *Circulation.* 2017;136:1331–45.
57. Thielmann M, Massoudy P, Jaeger BR, Neuhäuser M, Marggraf G, Sack S, Erbel R, Jakob H. Emergency re-revascularization with percutaneous coronary intervention, reoperation, or conservative treatment in patients with acute perioperative graft failure following coronary artery bypass surgery. *Eur J Cardiothorac Surg.* 2006;30:117–25.

58. Laflamme M, DeMey N, Bouchard D, Carrier M, Demers P, Pellerin M, Couture P, Perrault LP. Management of early postoperative coronary artery bypass graft failure. *Interact Cardiovasc Thorac Surg.* 2012;14:452–6.
59. Thielmann M, Massoudy P, Jaeger B, M Neuhäuser, G Marggraf, S Sack, R Erbel, H Jakob. Emergency re-revascularization with percutaneous coronary intervention, reoperation, or conservative treatment in patients with acute perioperative graft failure following coronary artery bypass surgery. *Eur J Cardio-Thoracic Surg.* 2006;30:117–25.
60. Rasmussen C, Thiis JJ, Clemmensen P, Efsen F, Arendrup H C, K Saunamäki, J K Madsen, G Pettersson. Significance and management of early graft failure after coronary artery bypass grafting. Feasibility and results of acute angiography and re-re-vascularization. *Eur J Cardio-Thoracic Surg.* 1997;12:847–52.
61. Davierwala PM, Verevkin A, Leontyev S, Misfeld M, Borger MA, Mohr FW. Impact of expeditious management of perioperative myocardial ischemia in patients undergoing isolated coronary artery bypass surgery. *Circulation.* 2013;128:S226-34.
62. Biancari F, Anttila V, Dell'Aquila AM, Airaksinen JKE, Brascia D. Control angiography for perioperative myocardial Ischemia after coronary surgery: meta-analysis. *J Cardiothorac Surg.* 2018;13:24.
63. Alqahtani F, Ziada KM, Badhwar V, Sandhu G, Rihal CS, Alkhouli M. Incidence, Predictors, and Outcomes of In-Hospital Percutaneous Coronary Intervention Following Coronary Artery Bypass Grafting. *J Am Coll Cardiol.* 2019;73:415–23.
64. Harskamp RE, Lopes RD, Baisden CE, de Winter RJ, Alexander JH. Saphenous vein graft failure after coronary artery bypass surgery: pathophysiology, management, and future directions. *Ann Surg.* 2013;257:824–33.
65. Brilakis ES, Edson R, Bhatt DL, Goldman S, Holmes DR, Rao S V, Shunk K, Rangan B V, Mavromatis K, Ramanathan K, Bavry AA, Garcia S, Latif F, Armstrong E, Jneid H, Conner TA, Wagner T, Karacsonyi J, Uyeda L, Ventura B, Alsleben A, Lu Y, Shih M-C, Banerjee S. Drug-eluting stents versus bare-metal stents in saphenous vein grafts: a double-blind, randomised trial. *Lancet.* 2018;391:1997–2007.

66. Mehilli J, Pache J, Abdel-Wahab M, Schulz S, Byrne RA, Tiroch K, Hausleiter J, Seyfarth M, Ott I, Ibrahim T, Fusaro M, Laugwitz K-L, Massberg S, Neumann F-J, Richardt G, Schömig A, Kastrati A. Drug-eluting versus bare-metal stents in saphenous vein graft lesions (ISAR-CABG): a randomised controlled superiority trial. *Lancet*. 2011;378:1071–8.
67. Colleran R, Kufner S, Mehilli J, Rosenbeiger C, Schüpke S, Hoppmann P, Joner M, Mankerious N, Fusaro M, Cassese S, Abdel-Wahab M, Neumann F-J, Richardt G, Ibrahim T, Schunkert H, Laugwitz K-L, Kastrati A, Byrne RA. Efficacy Over Time With Drug-Eluting Stents in Saphenous Vein Graft Lesions. *J Am Coll Cardiol*. 2018;71:1973–82.
68. Garg A, Rout A, Tayal R, Sharma A, Agrawal S, Kostis JB, Cohen M, Sharma S, Wasty N. Drug-eluting Stents Versus Bare-metal Stents for Saphenous Vein Graft Interventions: A Systematic Review and Meta-analysis of Studies With Longer Follow-up. *Curr Probl Cardiol*. 2019.
69. Morrison DA, Sethi G, Sacks J, Henderson WG, Grover F, Sedlis S, Esposito R. Percutaneous coronary intervention versus repeat bypass surgery for patients with medically refractory myocardial ischemia: AWESOME randomized trial and registry experience with post-CABG patients. *J Am Coll Cardiol*. 2002;40:1951–4.
70. Brilakis ES, Rao S V, Banerjee S, Goldman S, Shunk KA, Holmes DR, Honeycutt E, Roe MT. Percutaneous coronary intervention in native arteries versus bypass grafts in prior coronary artery bypass grafting patients: a report from the National Cardiovascular Data Registry. *JACC Cardiovasc Interv*. 2011;4:844–50.
71. Naidu SS, Turco MA, Mauri L, Coolong A, Popma JJ, Kereiakes DJ. Contemporary incidence and predictors of major adverse cardiac events after saphenous vein graft intervention with embolic protection (an AMEthyst trial substudy). *Am J Cardiol*. 2010;105:1060–4.
72. Baim DS, Wahr D, George B, Leon MB, Greenberg J, Cutlip DE, Kaya U, Popma JJ, Ho KKL, Kuntz RE. Randomized trial of a distal embolic protection device during percutaneous intervention of saphenous vein aorto-coronary bypass grafts. *Circulation*. 2002;105:1285–90.



73. Dixon SR, Mann JT, Lauer MA, Casale PN, Dippel EJ, Strumpf RK, Feldman RL, Shear W, Resar JR, Zimmer SD, O'Neill WW. A randomized, controlled trial of saphenous vein graft intervention with a filter-based distal embolic protection device: TRAP trial. *J Interv Cardiol.* 2005;18:233–41.
74. Paul TK, Bhatheja S, Panchal HB, Zheng S, Banerjee S, Rao S V, Guzman L, Beohar N, Zhao D, Mehran R, Mukherjee D. Outcomes of Saphenous Vein Graft Intervention With and Without Embolic Protection Device: A Comprehensive Review and Meta-Analysis. *Circ Cardiovasc Interv.* 2017;10.
75. Brilakis ES, O'Donnell CI, Penny W, Armstrong EJ, Tsai T, Maddox TM, Plomondon ME, Banerjee S, Rao S V, Garcia S, Nallamothu B, Shunk KA, Mavromatis K, Grunwald GK, Bhatt DL. Percutaneous Coronary Intervention in Native Coronary Arteries Versus Bypass Grafts in Patients With Prior Coronary Artery Bypass Graft Surgery: Insights From the Veterans Affairs Clinical Assessment, Reporting, and Tracking Program. *JACC Cardiovasc Interv.* 2016;9:884–93.
76. Wang N, Fulcher J, Abeysuriya N, Adams M, Lal S. Predictors of successful chronic total occlusion percutaneous coronary interventions: a systematic review and meta-analysis. *Heart.* 2018;104:517–24.
77. Stone GW, Maehara A, Lansky AJ, de Bruyne B, Cristea E, Mintz GS, Mehran R, McPherson J, Farhat N, Marso SP, Parise H, Templin B, White R, Zhang Z, Serruys PW. A Prospective Natural-History Study of Coronary Atherosclerosis. *N Engl J Med.* 2011;364:226–35.
78. Taniwaki M, Windecker S, Zaugg S, Stefanini GG, Baumgartner S, Zanchin T, Wenaweser P, Meier B, Ra L, Ju P. The association between in-stent neoatherosclerosis and native coronary artery disease progression: a long-term angiographic and optical coherence tomography cohort study. *Eur Heart J.* 2015;36(32):2167-76.
79. Cashin WL, Sanmarco ME, Nessim SA, Blankenhorn DH. Accelerated Progression of Atherosclerosis in Coronary Vessels with Minimal Lesions That Are Bypassed. *N Engl J Med.* 1984;311:824–8.
80. Pereg D, Fefer P, Samuel M, Wolff R, Czarnecki A, Deb S, Sparkes JD, Fremes SE, Strauss BH. Native Coronary Artery Patency After Coronary Artery Bypass Surgery. *JACC Cardiovasc Interv.* 2014;7:761–7.
81. Valgimigli M, Bueno H, Byrne RA, Collet J-P, Costa F, Jeppsson A, Jüni P, Kastrati A, Kolh P, Mauri L, Montalescot G, Neumann F-J, Petricevic M, Roffi M, Steg PG, Windecker S, Zamorano JL, Levine GN,

Badimon L, ESC Scientific Document Group; ESC Committee for Practice Guidelines (CPG); ESC National Cardiac Societies. 2017 ESC focused update on dual antiplatelet therapy in coronary artery disease developed in collaboration with EACTS. *Eur Heart J.* 2018;39:213–60.

82. Knuuti J, Wijns W, Saraste A, Capodanno D, Barbato E, Funck-Brentano C, E Prescott E, F Storey R, Deaton C, Cuisset T, Agewall S, Dickstein K, Edvardsen T, Escaned J, J Gersh B, Svitil P, Gilard M, Hasdai D, Hatala R, Mahfoud F, Masip J, Muneretto C, Valgimigli M, Achenbach S, J Bax J, ESC Scientific Document Group. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes The Task Force for the diagnosis and management of chronic. *Eur Heart J.* 2020;41(3):407–77.

83. Mäkikallio T, Holm NR, Lindsay M, Spence MS, Erglis A, Menown IBA, Trovik T, Eskola M, Romppanen H, Kellerth T, Ravkilde J, Jensen LO, Kalinauskas G, Linder RBA, Pentikainen M, Hervold A, Banning A, Zaman A, Cotton J, Eriksen E, Margus S, Sørensen HT, Nielsen PH, Niemelä M, Kervinen K, Lassen JF, Maeng M, Oldroyd K, Berg G, Walsh SJ, Hanratty CG, Kumsars I, Stradins P, Steigen TK, Fröbert O, Graham ANJ, Endresen PC, Corbascio M, Kajander O, Trivedi U, Hartikainen J, Anttila V, Hildick-Smith D, Thuesen L, Christiansen EH. Percutaneous coronary angioplasty versus coronary artery bypass grafting in treatment of unprotected left main stenosis (NOBLE): a prospective, randomised, open-label, non-inferiority trial. *Lancet.* 2016;388:2743–52.

84. Stone GW, Sabik JF, Serruys PW, Simonton CA, Généreux P, Puskas J, Kandzari DE, Morice M-C, Lembo N, Brown WM, Taggart DP, Banning A, Merkely B, Horkay F, Boonstra PW, van Boven AJ, Ungi I, Bogáts G, Mansour S, Noiseux N, Sabaté M, Pomar J, Hickey M, Gershlick A, Buszman P, Bochenek A, Schampaert E, Pagé P, Dressler O, Kosmidou I, Mehran R, Pocock SJ, Kappetein AP. Everolimus-Eluting Stents or Bypass Surgery for Left Main Coronary Artery Disease. *N Engl J Med.* 2016;375:2223–35.

85. Kandzari DE, Mauri L, Koolen JJ, Massaro JM, Doros G, Garcia-Garcia HM, Bennett J, Roguin A, Gharib EG, Cutlip DE, Waksman R. Ultrathin, bioresorbable polymer sirolimus-eluting stents versus thin, durable polymer everolimus-eluting stents in patients undergoing coronary revascularisation (BIOFLOW V): a randomised trial. *Lancet.* 2017;390:1843–52.

86. Biancari F, Dalén M, Ruggieri VG, Demal T, Gatti G, Onorati F, Faggian G, Rubino AS, Maselli D, Gherli R, Salsano A, Saccocci M, Santarpino G, Nicolini F, Tauriainen T, De Feo M, Airaksinen J, Rosato S, Perrotti A, Mariscalco G. Prognostic Impact of Multiple Prior Percutaneous Coronary Interventions in Patients Undergoing Coronary Artery Bypass Grafting. *J Am Heart Assoc.* 2018;7:e010089.
87. Escaned J. Secondary revascularization after CABG surgery. *Nat Rev Cardiol.* 2012;9:540–9.
88. Nashef SAM, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, Lockowandt U. EuroSCORE II. *Eur J Cardio-Thoracic Surg.* 2012;41:734–45.
89. Brener SJ, Lytle BW, Casserly IP, Ellis SG, Topol EJ, Lauer MS. Predictors of revascularization method and long-term outcome of percutaneous coronary intervention or repeat coronary bypass surgery in patients with multivessel coronary disease and previous coronary bypass surgery. *Eur Heart J.* 2006;27:413–8.
90. Serruys P W, Silber S, Garg S, Jan van Geuns R, Richardt G, Buszman PE, Ph D, Kelbæk H, Boven AJ Van, Ph D, Hofma SH, Ph D, Bressers M, Sc M, Gobbens P, Sc B, Negoita M. Comparison of Zotarolimus-Eluting and Everolimus-Eluting Coronary Stents. *N Engl J Med.* 2010;363(2):136-46.
91. Silber PS, Windecker PS, Vranckx P, Serruys P W, All R. Unrestricted randomised use of two new generation drug-eluting coronary stents: 2-year patient-related versus stent-related outcomes from the RESOLUTE All Comers trial. *Lancet.* 2011;377:1241–7.
92. Birgelen C Von, Sen H, Lam MK, Danse PW, Jessurun GAJ, Hautvast RWM, Houwelingen GK Van, R Schramm A , R Joe Gin MT, W Louwerenburg J , Frits H A F de Man , G Stoel M , M Löwik M , C M Linssen G , A M Saïd S , B Nienhuis M , M J Verhorst P , W Z Basalus M , J M Doggen C , Tandjung K. Third-generation zotarolimus-eluting and everolimus-eluting stents in all-comer patients requiring a percutaneous coronary intervention (DUTCH PEERS ): a randomised , single-blind , multicentre, non-inferiority trial. *Lancet.* 2014;383(9915):413-23.
93. Sen H , Kai Lam M , M Löwik M , W Danse P , A J Jessurun G , K Gert van Houwelingen , Antonio R L , R Melvyn Tjon Joe Gin , Raymond W M Hautvast , J Hans W Louwerenburg , Frits H A F de Man , G Stoel M , Liefke C van der Heijden , Gerard C M Linssen , Maarten J IJzerman, Kenneth Tandjung , Carine J

M Doggen, Clemens von Birgelen. Clinical Events and Patient-Reported Chest Pain in All-Comers Treated With Resolute Integrity and Promus Element Stents: 2-Year Follow-Up of the DUTCH PEERS (DUrable Polymer-Based STent CHallenge of Promus ElemEnt Versus ReSolute Integrity) Randomized Trial (TWENTE II). JACC Cardiovasc Interv.2015. 8(7):889-99.

94. Zocca P , Kok M M, Tandjung K , Danse PW , Jessurun G A J, Raymond W M Hautvast , van Houwelingen K , G Stoel M , R Schramm A , R Melvyn Tjon Joe Gin , Frits H A F de Man, Hartmann M , J Hans W Louwerenburg , Gerard C M Linssen, M Löwik M , J M Doggen C , von Birgelen C. 5-Year Outcome Following Randomized Treatment of All-Comers With Zotarolimus-Eluting Resolute Integrity and Everolimus-Eluting PROMUS Element Coronary Stents: Final Report of the DUTCH PEERS (TWENTE II) Trial 2018;11. JACC Cardiovasc Interv.2018. 11(5):462-469.

95. Saito S, Valdes-chavarri M, Richardt G, Moreno R, Romo I, Barbato E, Carrie D, Ando K, Merkely B, Kornowski R, Eltchaninoff H, James S, Wijns W. A Randomized, Prospective, Intercontinental Evaluation of a Bioresorbable Polymer Sirolimus-Eluting Coronary Stent System: The CENTURY II (Clinical Evaluation of New Terumo Drug-Eluting Coronary Stent System in the Treatment of Patients With Coronary Artery Disease) Trial Eur Heart J.2014;35(30):2021-31.

96. Wijns W, Valdes-chavarri M, Richardt G, Moreno R, Íñiguez-Romo A, Barbat oE, Carriè D, Ando K, Merkely B, Kornowski R, Eltchaninoff H, Stojkovic S, Saito S. Long-term clinical outcomes after bioresorbable and permanent polymer drug-eluting stent implantation : final five-year results of the CENTURY II randomised clinical trial. EuroIntervention 2018;14(3):e343-e351.

97. Pilgrim T, Heg D, Roffi M, Tüller D, Muller O, Vuilliomenet A, Cook S, Weilenmann D, Kaiser C, Jamshidi P, Fahrni T, Moschovitis A, Noble S, Eberli R F, Wenaweser P, Juni P, Windecker S. Ultrathin strut biodegradable polymer sirolimus-eluting stent versus durable polymer everolimus-eluting stent for percutaneous coronary revascularisation (BIOSCIENCE): a randomised, single-blind, non-inferiority trial. Lancet. 2014;384(9960):2111–22.

98. Zbinden R, Piccolo R, Heg D, Roffi M, Kurz DJ, Muller O, Vuilliomenet A, Cook S, Weilenmann D, Kaiser C, Jamshidi P, Franzone A, Eberli F, Peter J, Windecker S, Pilgrim T. Ultrathin Strut Biodegradable Polymer

*Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal*

Sirolimus-Eluting Stent Versus Durable-Polymer Everolimus-Eluting Stent for Percutaneous Coronary Revascularization: 2-Year Results of the BIOSCIENCE Trial. *J Am Heart Assoc.* 2016;5(3):e003255.

99. Pilgrim T, Piccolo R, Heg D, Roffi M, Tüller D, Müller O, Moarof I, Siontis GCM, Cook S, Weilenmann D, Kaiser C, Cuculi F, Hunziker L, Eberli FR, Jüni P, Windecker S. Ultrathin-strut, biodegradable-polymer, sirolimus-eluting stents versus thin-strut, durable-polymer, everolimus-eluting stents for percutaneous coronary revascularisation: 5-year outcomes of the BIOSCIENCE randomised trial. *Lancet.* 2018;392:737–46.

100. von Birgelen C, Kok MM, van der Heijden LC, Danse PW, Schotborgh CE, Scholte M, Gin RMTJ, Somi S, van Houwelingen KG, Stoel MG, de Man FHAF, Louwerenburg J (Hans) W, Hartmann M, Zocca P, Linssen GCM, van der Palen J, Doggen CJM, Löwik MM. Very thin strut biodegradable polymer everolimus-eluting and sirolimus-eluting stents versus durable polymer zotarolimus-eluting stents in allcomers with coronary artery disease (BIO-RESORT): a three-arm, randomised, non-inferiority trial. *Lancet.* 2016;388:2607–17.

101. Kok MM, Zocca P, Buiten RA, Danse PW, Schotborgh CE, Scholte M, Hartmann M, Stoel MG, van Houwelingen G, Linssen GCM, Doggen CJM, von Birgelen C. Two-year clinical outcome of all-comers treated with three highly dissimilar contemporary coronary drug-eluting stents in the randomised BIO-RESORT trial. *EuroIntervention.* 2018;14:915–23.

102. Buiten RA, Ploumen EH, Zocca P, Doggen CJM, Danse PW, Schotborgh CE, Scholte M, Stoel MG, Hartmann M, Tjon RM, Gin J, Somi S, Linssen GCM, Kok MM. Thin, Very Thin, or Ultrathin Strut Biodegradable- or Durable-Polymer- Coated Drug-Eluting Stents: 3-Year Outcomes of BIO-RESORT. *JACC Cardiovasc Interv.* 2019 Sep 9;12(17):1650-1660.

103. Jensen LO, Thayssen P, Maeng M, Junker A, Terkelsen CJ, Tilsted H, Hansen KN, Aaroe J, Kristensen SD, Hansen HS, Jensen SE, Madsen M, Bøtker HE. Randomized Comparison of a Biodegradable Polymer Ultrathin Strut Sirolimus-Eluting Stent With a Biodegradable Polymer Biolimus-Eluting Stent in Patients Treated With Percutaneous Coronary Intervention: The SOURT OUT VII Trial. *Circ Cardiovasc Interv.*

2016;9(7). 104. Jensen LO, Maeng M, Raungaard B, Hansen KN, Kahlert J, Jensen SE, Hansen HS,

Lassen JF, Bøtker HE. Two-year outcome after biodegradable polymer sirolimus- and biolimus-eluting coronary stents (from the randomised SORT OUT VII trial). *EuroIntervention*. 2018;13(13):1587–90.

105. Kandzari DE, Smits PC, Love MP, Ben-Yehuda O, Banai S, Robinson SD, Jonas M, Kornowski R, Bagur R, Iñiguez A, Danenberg H, Feldman R, Jauhar R, Chandna H, Parikh M, Y Perlman G, Balcells M, Markham P, Ozgu Ozan M, Genereux P, R Edelman E, B Leon M, W Stone G. Randomized Comparison of Ridaforolimus-Eluting and Zotarolimus-Eluting Coronary Stents in Patients with Coronary Artery Disease: Primary Results from the BIONICS Trial (BioNIR Ridaforolimus-Eluting Coronary Stent System in Coronary Stenosis). *Circulation* 2017;136(14):1304-1314.

106. von Birgelen C, Zocca P, Buiten RA, Jessurun GAJ, Schotborgh CE, Roguin A, Danse PW, Benit E, Aminian A, van Houwelingen KG, Anthonio RL, Stoel MG, Somi S, Hartmann M, Linssen GCM, Doggen CJM, Kok MM. Thin composite wire strut, durable polymer-coated (Resolute Onyx) versus ultrathin cobalt–chromium strut, bioresorbable polymer-coated (Orsiro) drug-eluting stents in allcomers with coronary artery disease (BIONYX): an international, single-blind, randomised non-inferiority trial. *Lancet*. 2018;392:1235–45.

107. Maeng M, Raungaard B, Kahlert J, Terkelsen CJ, Kristensen SD, Carstensen S, Jensen SE, Villadsen AB, Lassen JF, Thim T, Eftekhari A, Veien KT, Junker A, Jensen LO. Everolimus-Eluting Versus Biolimus-Eluting Stents With Biodegradable Polymers in Unselected Patients Undergoing Percutaneous Coronary Intervention: A Randomized Noninferiority Trial With 1-Year Follow-Up (SORT OUT VIII Trial). *JACC Cardiovasc Interv*. 2019;12(7):624-633.

108. Geisler T, Zürn C, Simonenko R, Rapin M, Kraibooj H, Kiliass A, Bigalke B, Stellos K, Schwab M, May AE, Herdeg C, Gawaz M. Early but not late stent thrombosis is influenced by residual platelet aggregation in patients undergoing coronary interventions. *Eur Heart J*. 2010;31:59–66.

109. Waksman R, Kirtane AJ, Torguson R, Cohen DJ, Ryan T, Räber L, Applegate R, Waxman S, Gordon P, Kaneshige K, Leon MB. Correlates and Outcomes of Late and Very Late Drug-Eluting Stent Thrombosis. *JACC Cardiovasc Interv*. 2014;7:1093–102.

110. Taniwaki M, Radu MD, Zaugg S, Amabile N, Garcia-Garcia HM, Yamaji K, Jørgensen E, Kelbæk H, Pilgrim T, Caussin C, Zanchin T, Veugeois A, Abildgaard U, Jüni P, Cook S, Koskinas KC, Windecker S, Räber L. Mechanisms of Very Late Drug-Eluting Stent Thrombosis Assessed by Optical Coherence Tomography. *Circulation*. 2016;133:650–60.
111. Cassese S, Byrne RA, Tada T, Piniček S, Joner M, Ibrahim T, King LA, Fusaro M, Laugwitz K-L, Kastrati A. Incidence and predictors of restenosis after coronary stenting in 10 004 patients with surveillance angiography. *Heart*. 2014;100:153–9.
112. Cayla G, Hulot JS, O'Connor SA, Pathak A, Scott SA, Gruel Y, Silvain J, Vignalou JB, Huerre Y, De La Briolle A, Allanic F, Beygui F, Barthélémy O, Montalescot G, Collet JP. Clinical, angiographic, and genetic factors associated with early coronary stent thrombosis. *JAMA*. 2011;306:1765–74.
113. Kuramitsu S, Ohya M, Shinozaki T, Otake H, Horie K, Kawamoto H, Yamanaka F, Natsuaki M, Shiomi H, Nakazawa G, Ando K, Kadota K, Saito S, Kimura T. Risk Factors and Long-Term Clinical Outcomes of Second-Generation Drug-Eluting Stent Thrombosis: Insights from the REAL-ST Registry. *Circ Cardiovasc Interv*. 2019;12:1–9.
114. Mehran R, Dangas G, Abizaid AS, Mintz GS, Lansky AJ, Satler LF, Pichard AD, Kent KM, Stone GW, Leon MB. Angiographic patterns of in-stent restenosis: classification and implications for long-term outcome. *Circulation*. 1999;100:1872–8.
115. Garcia-Guimaraes M, Antuña P, Maruri-Sanchez R, Vera A, Cuesta J, Bastante T, Rivero F, Alfonso F. Calcified neoatherosclerosis causing in-stent restenosis. *Coron Artery Dis*. 2019;30:1–8.
116. Joner M, Koppa T, Byrne RA, Castellanos MI, Lewerich J, Novotny J, Guagliumi G, Xhepa E, Adriaenssens T, Godschalk TC, Malik N, Alfonso F, Tada T, Neumann F-J, Desmet W, ten Berg JM, Gershlick AH, Feldman LJ, Massberg S, Kastrati A. Neoatherosclerosis in Patients With Coronary Stent Thrombosis. *JACC Cardiovasc Interv*. 2018;11:1340–50.
117. Xhepa E, Byrne RA, Rivero F, Rroku A, Cuesta J, Ndrepepa G, Kufner S, Valiente TB, Cassese S, Garcia-Guimaraes M, Lahmann AL, Rai H, Schunkert H, Joner M, Pérez-Vizcayno MJ, Gonzalo N, Alfonso F,

Kastrati A. Qualitative and quantitative neointimal characterization by optical coherence tomography in patients presenting with in-stent restenosis. Clin Res Cardiol. 2019;108(9):1059-1068.

Copyright EuroIntervention